

INDOXACARB GROUP 22A INSECTICIDE

Cameo SC

ACTIVE INGREDIENT:

Indoxacarb: (S)-methyl 7-chloro-2,5-dihydro-2-[[[(methoxycarbonyl)[4-(trifluoromethoxy)phenyl]amino]carbonyl]indeno[1,2-e][1,3,4]oxadiazine-4a(3H)-carboxylate. 14.5%

OTHER INGREDIENTS: 85.5%

TOTAL: **100.0%**

Contains 1.25 lbs. indoxacarb per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

**See label booklet for complete First Aid, Precautionary Statements,
Directions For Use, and Storage and Disposal.**

Manufactured For:

Sharda USA LLC 

7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

EPA Reg. No. 83529-154

EPA Est. No. **SC** 39578-TX-001; **VP** 07401-TX-001; **MA** 83411-MN-001;
AF 86555-MO-001; **CS** 70815-GA-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Contents: 1 Gallon

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222.</p> <p>For general information about this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Harmful if inhaled. Avoid breathing (dust, vapor, or spray mist). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse and before eating, drinking, chewing gum, using tobacco, or using the toilet. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Water-proof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, birds, fish, and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment wash water or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 24 hours. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems. Do not apply to any impervious surfaces which may contact or lead directly to surface water, storm drains, or urban runoff conveyance systems (gutters).

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging in the treatment area.

For fields to which applications of **Cameo SC** will be made, construct a vegetative filter strip if one does not already exist. Existing and new filter strips must be, at a minimum, 10-foot-wide and composed of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds). Vegetative filter strips must be maintained to optimize their utility. Only apply products containing indoxacarb onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This product can only be used in accordance with the Directions for Use on this label. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Water-proof gloves
- Shoes plus socks

PRODUCT INFORMATION

Cameo SC is a suspension concentrate that can be applied as a foliar spray to control many important insects. **Cameo SC** is mixed with water for application.

Always shake **Cameo SC** well before use.

Use Restrictions

- Do not use in greenhouses.
- **Chemigation:** Refer to supplemental or Special Local Need (SLN) labeling for use directions for chemigation. Do not apply this product through any irrigation system unless the supplemental or SLN labeling on chemigation is followed.

INTEGRATED PEST MANAGEMENT

Sharda USA LLC supports the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an IPM program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, rotation of insecticides with different modes-of-action, and treating when target pest populations reach locally determined action thresholds. Consult your State Cooperative Extension Service, professional consultants, or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop or site systems in your area.

BENEFICIAL ARTHROPODS

Other than reducing the target pest species as a food source, **Cameo SC** helps conserve certain beneficial arthropods, including parasitic wasps, predatory mites, big-eyed bugs, damsel bugs, minute pirate bugs, and spiders. While these beneficials cannot be relied upon to control pests, they are of potential value and can be monitored along with pests in pest management programs on these crops.

SCOUTING

Monitor insect populations to determine whether or not there is a need for application of **Cameo SC** based on locally determined economic thresholds. More than one treatment of **Cameo SC** may be required to control a population of pests. Refer to **CROP USE DIRECTIONS** table for maximum application rate restrictions for each crop.

INSECT RESISTANCE MANAGEMENT

For resistance management, **Cameo SC** contains Indoxacarb and is classified in the oxadiazine chemical class as a Group 22A insecticide, voltage-dependent sodium channel blockers.

Any insect population may contain individuals naturally resistant to **Cameo SC** and other Group 22A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies must be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of **Cameo SC** or other Group 22A insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues for the targeted pests between the individual components of a mixture.
- In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures must be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological, and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance, contact Sharda USA LLC or representative.

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications:

- Do not release spray at a height greater than 10 ft. above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 mph at the application site. If wind speed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed-wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Applicators must use 1/2 swath displacement upwind at the downwind edge of field.
- Do not apply during temperature inversions.

Ground Boom Applications:

- Users must only apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy unless making a turf, pasture, or rangeland application, in which case applicators may apply with a nozzle height no more than 4 feet above the ground.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size - Aircraft

- **Adjust Nozzles** - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles must be oriented parallel with the airflow in flight.

BOOM HEIGHT - Ground Boom

For ground equipment, the boom must remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.** Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

AIR ASSISTED (AIR BLAST) - FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring. Note: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Refer to the **APPLICATION INSTRUCTION** section of this label to determine if use of an air assisted sprayer is recommended.

APPLICATION INSTRUCTIONS

Apply at the specified rates when insect populations reach locally determined economic thresholds. Consult the State Cooperative Extension Service, professional consultants, or other qualified authorities to determine appropriate threshold levels for treatment in your area.

Cameo SC applications should target eggs and small instar larvae.

Follow-up treatments of **Cameo SC** should be applied, as needed, to keep pest populations within threshold limits. **Cameo SC** should be applied at 5- to 7-day intervals or as needed to manage specific target insect pests, as specified in the specific crop sections to maintain control.

Use sufficient water to obtain thorough, uniform coverage.

For aerial application, use a minimum of 3 gals. water per acre (gpa) (minimum of 5 gals. water per acre in Arizona and California). For ground application, use a minimum of 5 gals. water per acre. Higher gallonage will provide better coverage and performance.

MIXING PROCEDURES

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Spray equipment must be clean and free of previous pesticide deposits before applying **Cameo SC**. Fill spray tank 1/4 - 1/2 full of water. Add **Cameo SC** directly to spray tank. Mix thoroughly to fully disperse the insecticide; once dispersed continued agitation is required. Use mechanical or hydraulic means; do not use air agitation. Spray mix must not be stored overnight in spray tank.

Compatibility - Since formulations may be changed and new ones introduced, it is recommended that users premix a small quantity of a desired tank mix and observe for possible adverse changes (settling out, flocculation, etc.). Avoid mixtures of several materials and very concentrated spray mixtures.

Tank Mixing Sequence - Add different formulation types in the sequence indicated below, unless otherwise specified by manufacturer directions for use or by local expertise. Allow time for complete mixing and dispersion after addition of each product.

1. Water soluble bags.
2. Water dispersible granules.
3. Wettable powders.
4. Water-based suspension concentrates.
5. Water soluble concentrates.
6. **Cameo SC** and other oil-based suspension concentrates.
7. Emulsifiable concentrates.
8. Adjuvants, surfactants, oils.
9. Soluble fertilizers.
10. Drift retardants.

SPRAY TANK CLEANOUT

Prior to application, start with clean, well-maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove.

Drain spray equipment. Thoroughly rinse sprayer and flush hoses, boom, and nozzles with clean water.

Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

CROP ROTATION

Crops that are on this label and broccoli, brussels sprouts, cabbage, cauliflower, Chinese broccoli, Chinese cabbage, Chinese mustard cabbage, kohlrabi, lettuce (head and leaf varieties), pepper (bell and non-bell), potato, sweet corn, and tomato may be planted immediately following harvest.

Restriction: Do not plant for food or feed any other crops not registered for use with indoxacarb for 30 days after last use.

CROP USE DIRECTIONS

ALFALFA

Pests	Cameo SC Rate per Acre	Acres Treated per Gal. of Cameo SC
Alfalfa Caterpillar Alfalfa Weevil (Larvae) Beet Armyworm Egyptian Alfalfa Weevil (Larvae) Granulate Cutworm	6.7 - 11.3 fl. oz. (0.065 - 0.11 lb. a.i.)	11.5 - 19
Potato Leafhopper (Except California) (Suppression Only) Western Yellowstriped Armyworm	9.2 - 11.3 fl. oz. (0.09 - 0.11 lb. a.i.)	11.5 - 14
Apply lower listed rates on light to moderate infestations. Use intermediate to high rates on heavier infestations or when later instar larvae exist. Use the highest listed rate for controlling severe infestations or when longer residual control is desired.		
Restrictions:		
<ul style="list-style-type: none">• Make no more than 1 application per cutting.• Do not apply more than 45 fl. oz. (0.44 lb. a.i.) of Cameo SC per acre per crop season.• Pre-Harvest Interval (PHI): 7 days. For alfalfa, harvest is defined as when the crop is cut.• Restricted-Entry Interval: 12 hours• Not for use in alfalfa grown for seed or in alfalfa grown for alfalfa sprouts for human consumption.		

PEANUT

Pests	Cameo SC Rate per Acre	Acres Treated per Gal. of Cameo SC
Beet Armyworm Corn Earworm Fall Armyworm Granulate Cutworm Rednecked Peanutworm (except California)	9.2 - 11.3 fl. oz. (0.09 - 0.11 lb. a.i.)	11.5 - 14
Restrictions: <ul style="list-style-type: none"> • Do not apply more than 45 fl. oz. (0.44 lb. a.i.) of Cameo SC per acre per crop. • Retreatment Interval: 5 days • Pre-Harvest Interval (PHI): 14 days • Restricted-Entry Interval: 12 hours 		

SOYBEAN (except California)

Pests	Cameo SC Rate per Acre	Acres Treated per Gal. of Cameo SC
Beet Armyworm Cabbage Looper Corn Earworm Fall Armyworm Green Cloverworm Soybean Looper Velvetbean Caterpillar (Suppression Only) Yellowstriped Armyworm	5.6 - 11.3 fl. oz. (0.055 - 0.11 lb. a.i.)	11.5 - 22.8
Restrictions: <ul style="list-style-type: none"> • Do not apply more than 45 fl. oz. (0.44 lb. a.i.) of Cameo SC per acre per crop. • Retreatment Interval: 5 days • Pre-Harvest Interval (PHI): 21 days • Restricted-Entry Interval: 12 hours • Do not feed or graze livestock on treated fields. 		

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Do not subject to temperatures below 32°F. Store product in original container only in a location inaccessible to children and pets. Not for use or storage in or around the home.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Less Than or Equal to 5 Gallons. Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures allowed by State and local authorities.

Greater Than 5 Gallons. Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by other procedures allowed by State and local authorities.

For Bulk and Mini-Bulk Containers. Refillable container. Refill this container with pesticide only. Do not use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by State and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and, to the extent consistent with applicable law, Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

All trademarks are the property of their respective owners.

INDOXACARB GROUP 22A INSECTICIDE

Cameo SC

ACTIVE INGREDIENT:	WT. BY %
Intoxcarb: (S)-methyl 7-chloro-2,5-dihydro-2-[[[methoxy(aryl)oxy]oxy]oxy]oxy-4a(3H)-carboxylate	14.5%
OTHER INGREDIENTS:	85.5%
TOTAL:	100.0%

Contains 1.25 lbs. Indoxcarb per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID - IF SWALLOWED: • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. **IF ON SKIN OR CLOTHING:** • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice. **IF IN EYES:** • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. **IF INHALED:** • Move person to fresh air. • If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER - Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222. For general information about this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PST, or at <http://npic.orst.edu>

See label booklet for complete Precautionary Statements and Directions For Use.

PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION - Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Harmful if inhaled. Avoid breathing (dust, vapor, or spray mist). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse and before eating, drinking, chewing gum, using tobacco, or using the toilet. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS - This pesticide is toxic to mammals, birds, fish, and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment wash water or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 24 hours. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems. Do not apply to any impervious surfaces which may contact or lead directly to surface water, storm drains, or urban runoff conveyance systems (gutters). This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging in the treatment area. For fields in which applications of **Cameo SC** will be made, construct a vegetative filter strip if one does not already exist. Existing and new filter strips must be, at a minimum, 100-foot-wide and composed of grasses or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds). Vegetative filter strips must be maintained to optimize their utility. Only apply products containing indoxcarb onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **PESTICIDE STORAGE:** Do not subject to temperatures below 32°F. Store product in original container only in a location inaccessible to children and pets. Not for use or storage in or around the home. **PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER HANDLING: Less Than or Equal to 5 Gallons.** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures allowed by State and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

Manufactured For:

Sharda USA LLC, 7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707
EPA Reg. No. 83529-154 EPA Est. No. SC 39578-TX-001; VP 07401-TX-001
MA 83411-MN-001; AF 86555-MO-001; CS 70165-GA-001

The EPA Establishment Number is identified by the circled letters above that match the first two letters in the batch number.

Net Contents: 1 Gallon

OPEN HERE