

DICAMBA GROUP 4 HERBICIDE

Disha DMA

Herbicide For Weed Control in Corn, Sorghum, Small Grains (wheat, barley, and oats), Pasture, Hay, Rangeland, General Farmstead (Non-Cropland), Rights-of-Way, Public Utility and Industrial Areas, Fallow, Cotton, Sugarcane, Asparagus, Turf, and Grass Seed Crops

ACTIVE INGREDIENT:	WT. BY %
Dimethylamine salt of dicamba (3, 6-dichloro-o-anisic acid).....	49.77%
OTHER INGREDIENTS:	50.23%
TOTAL:	100.00%

Contains 41.71% 3, 6-dichloro-o-anisic acid (dicamba) or 4 pounds per gallon (480 g/L).

**KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

See label booklet for complete Precautionary Statements and Directions For Use.

EPA Reg. No.: 83529-110

EPA Est. No.:

- 70815-GA-002
- 44616-MO-002
- 39578-TX-001
- 93796-GRC-001

Manufactured For:

Sharda USA LLC 

7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

Net Contents: 265 Gallons

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at **1-800-222-1222**.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a well-ventilated area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING [Less Than 5 Gallons]: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONTAINER HANDLING [Greater Than 5 Gallons]: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

CONTAINER HANDLING [For Bulk and Mini-Bulk Containers]: Refillable container. Refill this container with pesticide only. Do not use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by State and local authorities.

OPEN HERE 

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at **1-800-222-1222**.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All mixers, loaders, and applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material (except for applicators using ground boom equipment, pilots and flaggers)
- Protective eyewear

See **ENGINEERING CONTROLS** for additional requirements and exceptions.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use cockpits in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [(40 CFR 170.240(d)(4-6))].

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling the product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Apply this product only as directed on label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **24 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls worn over short-sleeve shirt and short pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material
- Chemical-resistant headgear for overhead exposure
- Protective eyewear

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, nurseries, or greenhouses.

DO NOT enter or allow others to enter the treated area until sprays have dried.

BEST STEWARDSHIP PRACTICES

Disha DMA provides effective broadleaf weed and brush control when properly applied. Best stewardship practices in all mixing, loading, and application operations not only maximize weed control, but also protect ground and surface waters and minimize off-target movement.

This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

GROUND AND SURFACE WATERS PROTECTION

Point Source Contamination: To prevent point source contamination, do not mix or load this pesticide product within 50 feet of wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. Do not apply pesticide product within 50 feet of wells. This setback does not apply to properly capped or plugged abandoned wells according to State and local requirements and does not apply to impervious pad or properly diked mixing/loading areas as described in this section. Mixing, loading, rinsing, or washing operations performed within 50 feet of a well are allowed only when conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be on or moved across the pad. The pad must be self-contained to prevent surface water flow over or from the pad. The pad capacity must be maintained at 110% that of the largest pesticide container or application equipment used on the pad and have sufficient capacity to contain all product spills, equipment or container leaks, equipment wash waters, and rainwater that may fall on the pad. The containment capacity does not apply to vehicles delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment. Care must be taken when using this product to prevent: a) back siphoning into wells, b) spills or c) improper disposal of excess pesticide, spray mixtures or rinsates. Check valves or anti-siphoning devices must be used on all mixing equipment.

Movement by surface runoff of through soil: Do not apply under conditions which favor runoff. Do not apply to impervious substrates such as paved or highly compacted surfaces in areas with high potential for ground water contamination. Ground water contamination may occur in areas where soils are permeable or coarse and ground water is near the surface. Do not apply to soils classified as sand with less than 3% organic matter and where ground water depth is shallow. To minimize the possibility of ground water contamination, carefully follow specified application rates as affected by soil type in the **PRODUCT INFORMATION** section of this label.

Movement by water erosion of treated soil: Do not apply or incorporate this product through any type of irrigation equipment nor by flood or furrow irrigation. Ensure treated areas have received at least ½ inch rainfall (or irrigation) before using tailwater for subsequent irrigation of other fields.

PRODUCT INFORMATION

Before applying **Disha DMA** read all directions and precautions appearing on the container label and in this booklet. Failure to follow all directions and precautions may result in unsatisfactory weed control, crop injury, or illegal residues.

The following directions apply to all uses of **Disha DMA**. Additional precautions and restrictions will be found in each specific use section.

Use Restrictions:

- Do not treat irrigation ditches or water used for crop irrigation or domestic uses.
- Do not apply this product through any type of irrigation system.
- Rates of use of this product must not exceed 2 pints per acre, per application; and 4 pints per acre, per year.
- Rates of use of dicamba acid equivalent must not exceed a maximum single application rate of 1.0 lb. per acre; and an annual application rate of 2.0 lbs. per acre. These restrictions apply to this product and possible mixtures with other products containing dicamba.

SENSITIVE CROP PRECAUTIONS

Disha DMA may cause injury to desirable trees and plants, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes, and other broadleaf plants when contacting their roots, stems, or foliage. These plants are most sensitive to **Disha DMA** during their development or growing stage.

FOLLOW THE PRECAUTIONS LISTED ON THIS LABEL WHEN USING DISHA DMA.

Use Precautions:

- Use coarse sprays to avoid potential herbicide drift. Select nozzles which are designed to produce minimal amounts of fine spray particles. Examples of nozzles designed to produce coarse sprays via ground applications are Delavan Raindrops, Spraying Systems XR flat fans or large capacity flood nozzles such as D10, TK10, or greater capacity tips. Keep the spray pressure at or below 20 psi and the spray volume at or above 20 gpa, unless otherwise required by the manufacturer of drift reducing nozzles. Consult with your spray nozzle supplier concerning the choice of drift-reducing nozzles.
- Agriculturally approved drift-reducing additives may be used.
- To avoid injury to desirable plants, thoroughly clean the equipment used to apply **Disha DMA** before using the equipment to apply any other chemical (refer to the **Procedure For Cleaning Spray Equipment** section).

Use Restrictions:

- Do not treat areas where either possible downward movement into the soil or surface washing may cause contact of **Disha DMA** with the roots of desirable plants such as trees and shrubs.
- Do not apply when spray particles maybe carried by air currents to areas where sensitive crops and plants are growing or when temperature inversions exist.
- Do not spray near sensitive plants if wind is gusty or in excess of 5 mph and moving in the direction of adjacent sensitive plants.
- Do not apply **Disha DMA** adjacent to sensitive crops when the temperature on the day of application is expected to exceed 85°F as drift is more likely to occur.

All crop uses of **Disha DMA** are intended for a normal growing interval between planting and harvest. No crop rotation restrictions exist if normal harvest of treated crop has occurred. If this interval is shortened, such as in cover crops that will be plowed under, do not follow up with the planting of a sensitive crop.

Crops growing under stress conditions such as drought, poor fertility, or foliar damage due to hail, wind or insects, can exhibit various injury symptoms that may be more pronounced if herbicides are applied. Consult your local or State authorities for possible application restrictions and advice concerning these and other special local use situations. Tank mix instructions are for use only in states where the tank mix product and application site are registered.

MIXING AND APPLICATION

UNLESS OTHERWISE SPECIFIED UNDER THE INDIVIDUAL USE HEADINGS OF THIS BOOKLET, THE FOLLOWING DIRECTIONS APPLY TO ALL CROP AND NON-CROP USES OF **DISHA DMA**. REFER TO INDIVIDUAL USE SECTIONS FOR ADDITIONAL PRECAUTIONS, RESTRICTIONS, APPLICATION RATES, AND TIMINGS.

Disha DMA is a water-soluble formulation that can be applied using water or sprayable fluid fertilizer as the carrier. If a fluid fertilizer is to be used, a compatibility test (refer to the **Compatibility Test** section) should be made prior to tank mixing.

Ground or aerial application equipment which will give good spray coverage of weed foliage should be used. HOWEVER, **DO NOT** USE AERIAL APPLICATION EQUIPMENT IF SPRAY PARTICLES CAN BE CARRIED BY WIND INTO AREAS WHERE SENSITIVE CROPS OR PLANTS ARE GROWING OR WHEN TEMPERATURE INVERSIONS EXIST.

Apply 3 to 50 gals. of diluted spray per treated acre when using ground application equipment or 1 - 10 gals. of diluted spray per treated acre (2 - 20 gals. of diluted spray per treated acre for pre-harvest uses) in a water-based carrier when using aerial application equipment. Use the higher level of the listed spray volumes when treating dense or tall vegetation. Use coarse sprays. Select nozzles designed to produce minimal amounts of fine spray particles. Spray with nozzles as close to the weeds as is practical for good weed coverage.

To avoid uneven spray coverage, **Disha DMA** should not be applied during periods of gusty wind or when wind is in excess of 15 mph.

Avoid disturbing (e.g., cultivating or mowing) treated areas for at least 7 days following application.

Band Treatment

Disha DMA may be applied as a band treatment. Use the following formulas to determine the appropriate rate and volume per treated acre:

$$\frac{\text{Band Width (Inches)}}{\text{Row Width (Inches)}} \times \text{Broadcast Rate per Acre} = \text{Band Rate per Acre}$$

$$\frac{\text{Band Width (Inches)}}{\text{Row Width (Inches)}} \times \text{Broadcast Volume per Acre} = \text{Band Volume per Acre}$$

Compatibility Test

Before mixing in the spray tank, it is advisable to test compatibility by mixing all components in a small container in proportionate quantities (refer to the following table).

Amount of Herbicide to Add to 1 Pint of Spray Carrier (Assuming volume is 25 gals. per acre)		
Herbicide Formulations	Rate per Acre	Level Teaspoons
Dry	1 lb.	1 ½
Liquid	1 pt.	½

If herbicide(s) do not ball-up or form flakes, sludge, gels, oily films or layers, or other precipitates, then the tested spray mix is compatible. Usually, incompatibility in any of the previously described forms will occur within 5 minutes after mixing.

If components are incompatible, the use of a compatibility agent is recommended. Rerun the compatibility test with a suitable compatibility agent (¼ teaspoon is equivalent to 2 pints per 100 gallons of fluid fertilizer).

Procedure For Cleaning Spray Equipment

The steps listed below are suggested for thorough cleaning of spray equipment following applications of **Disha DMA** or tank mixes of **Disha DMA** or tank mixes of **Disha DMA** plus 2,4-D amine:

- Hose down thoroughly the inside as well as outside surfaces of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of the rinse water.
- Fill tank with water while adding 1 quart of household ammonia for every 25 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer system for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
- Flush the solution out of the spray tank through the boom.
- Remove the nozzles and screens and flush the system with 2 full tanks of water.

Steps 5 through 8 are suggested for thorough cleaning of spray equipment used to apply **Disha DMA** as a tank mix with wettable powders (WP), emulsifiable concentrates (EC), or other types of water-dispersible formulations. **Disha DMA** tank mixes with water-dispersible formulations require the use of a water/detergent rinse.

- Complete step 1.
- Fill tank with water while adding 2 lbs. of detergent for every 40 gals. of water. Operate the pump to circulate the detergent solution through the sprayer system for 5 to 10 minutes and discharge a small amount of the solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
- Flush the detergent solution out of the spray tank through the boom.
- Repeat step 1, and follow with steps 2, 3, and 4.

WEED LIST

This is a list of weeds which may be treated with **Disha DMA** in accordance with this label as specified under the **CROP-SPECIFIC INFORMATION** sections of the individual Use headings. Proper usage of this product will give control or growth suppression of many annual, biennial, and perennial broadleaf weeds, and many woody brush and vine species including:

ANNUALS		
Amaranth, Spiny (Spiny Pigweed)	Kochia	Radish, Wild
Aster, Slender	Ladysthumb	Ragweed, Common
Bedstraw	Lambsquarters, Common	Ragweed, Giant (Buffaloweed)
Beggarweed, Florida	Lambsquarters (Triazine-Resistant)	Ragweed, Lance-Leaf
Broomweed, Common	Lettuce, Prickly	Rubberweed, Bitter (Bitterweed)
Buckwheat, Wild	Mallow, Common	Sesbania, Hemp
Buffalobur	Mallow, Venice	Shepherd's Purse
Burclover, California	Mayweed	Sicklepod

(continued)

ANNUALS (cont.)		
Burcucumber	Morningglory, Ivyleaf	Sida, Prickly (Teaweed)
Buttercup, Roughseed	Morningglory, Tall	Smartweed, Green
Carpetweed	Mustard, Tansy	Smartweed, Pennsylvania
Catchfly, Nightflowering	Mustard, Wild	Sneezeweed, Bitter
Chamomile, Corn	Mustard (Yellowtops)	Sowthistle, Annual
Chickweed, Common	Nightshade, Black	Sowthistle, Spiny
Clovers (Annual)	Pennycress, Field (Fanweed Frenchweed, Stinkweed)	Spikeweed, Common
Cockle, Corn	Pepperweed, Virginia (Peppergrass)	Spurge, Prostrate
Cockle, Cow	Pigweed, Prostrate	Spurry, Corn
Cocklebur, Common	Pigweed, Redroot (Carelessweed)	Starbur, Bristly
Croton, Tropic	Pigweed, Rough	Sumpweed, Rough
Croton, Woolly	Pigweed, Smooth	Sunflower, Common (Wild)
Daisy, English	Pigweed (Triazine-Resistant)	Sunflower, Volunteer
Evening Primrose, Cutleaf	Pigweed, Tumble	Thistle, Russian
Fleabane, Annual	Poorjoe	Velvetleaf
Goosefoot, Nettleleaf	Puncturevine	Waterhemp
Henbit	Purslane, Common	Water Primrose, Winged
Jimsonweed	Pusley, Florida	Wormwood, Annual
Knotweed		
BIENNIALS		
Burdock, Common	Knapweed, Diffuse	Sweetclover
Carrot, Wild (Queen Anne's Lace)	Knapweed, Spotted	Teasel
Cockle, White	Mallow, Dwarf	Thistle, Bull
Evening Primrose, Common	Plantain, Bracted	Thistle, Milk
Geranium, Carolina	Ragwort, Tansy	Thistle, Musk
Gromwell	Starthistle, Yellow	Thistle, Plumelless
PERENNIALS		
Alfalfa ¹	Fern, Bracken	Redvine
Artichoke, Jerusalem	Garlic, Wild	Sericia Lespedeza
Aster, Spiny	Goldenrod, Canada	Smartweed, Swamp
Aster, Whiteheath	Goldenrod, Missouri	Snakeweed, Broom
Beadstraw, Smooth	Goldenweed, Common	Sorrel, Red (Sheep Sorrel) ¹
Bindweed, Field	Hawkweed	Sowthistle
Bindweed, Hedge	Horsenettle, Carolina	Sowthistle, Perennial
Blueweed, Texas	Ironweed	Spurge, Leafy
Bursage (Bur Ragweed Lakewood, Povertyweed) ¹	Knapweed, Black Knapweed, Russian	Sundrop, Halfshrub (Evening Primrose) Thistle, Canada
Bursage, Woollyleaf (Lakewood)	Mare's Tail (Horseweed)	Toadflax, Dalmation
Buttercup, Tall	Milkweed, Climbing	Tropical Soda Apple
Campion, Bladder	Milkweed, Common	Trumpetcreeper (Buckvine)
Chickweed, Field	Milkweed, Honeyvine	Vetch
Chickweed (Mouseear, Canada)	Milkweed, Western Whorled	Waterhemlock
Chicory	Nettle, Stinging	Waterprimrose, Creeping
Clover, Hop ¹	Nightshade, Silverleaf (White Horsenettle)	Woodsorrel, Creeping (Common Yellow) ¹
Dandelion, Common ¹	Onion, Wild	Wormwood, Common
Dock, Broadleaf (Bitterdock) ¹	Plantain, Broadleaf ¹	Wormwood, Louisiana
Dock, Curly ¹	Plantain, Buckhorn	Yankeeeweed ¹
Dogbane, Hemp	Pokeweed	Yarrow, Common
Dogfennel (Cypressweed) ¹	Ragweed, Western	
WOODY		
Alder	Hawthorn (Thornapple)*	Plum, Sand (Wild Plum)*
Ash	Hemlock	Rabbitbrush
Aspen	Hickory	Red Cedar, Eastern*
Basswood	Honeylocust	Rose, McCartney*
Beech	Honeysuckle	Rose, Multiflora*
¹ Noted perennials may be controlled using Disha DMA at rates lower than those directed for other listed perennial weeds.		
*Growth Suppression.		

WOODY (cont.)		
Birch	Hornbeam	Sagebrush, Fringe
Blackberry*	Huckleberry	Sassafras
Blackgum*	Huisache	Serviceberry
Cedar*	Ivy, Poison	Spicebush
Cherry	Kudzu	Spruce
Chinquapin	Locust, Black	Sumac
Cottonwood	Maple	Sweetgum*
Creosotebush*	Mesquite	Sycamore
Cucumbertree	Oak	Tarbrush
Dewberry*	Oak, Poison	Willow
Dogwood*	Olive, Russian	Witchhazel
Elm	Persimmon, Eastern	Yaupon*
Grape	Pine	Yucca*

¹Noted perennials may be controlled using **Disha DMA** at rates lower than those directed for other listed perennial weeds.
^{*}Growth Suppression.

CROP-SPECIFIC INFORMATION

ASPARAGUS

Apply **Disha DMA** to emerged and actively growing weeds in 40 - 60 gals. of diluted spray per treated acre immediately after cutting the field, but at least 24 hours before the next cutting. If spray contacts emerged spears, crooking (twisting) of some spears may result. If such crooking occurs, discard affected spears.

Note: Observe all precautions. Read and follow mixing, application, and cleaning instructions.

Restrictions:

- Multiple applications may be made per growing season. Do not exceed a total of 1 pt. of **Disha DMA** (0.5 lb. a.e. dicamba) per treated acre per crop year.
- Do not harvest prior to 24 hours after treatment.
- Do not use in the Coachella Valley of California.
- **Disha DMA** contains 0.5 lb. a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 0.5 lb. of a.e. per acre per application.

Weeds	Rate per Treated Acre
Mustard, Black, Pigweed, Redroot (Carelessweed) Sowthistle, Annual Thistle, Canada ¹ Thistle, Russian	½ - 1.0 pt. (¼ - ½ lb. a.e.)
Bindweed, Field ¹ Chickweed, Common Goosefoot, Nettleleaf Radish, Wild Thistle, Milk	1.0 pt. (½ lb. a.e.)

¹**Disha DMA** may be applied in a tank mixture with either 2,4-D or Roundup herbicide for improved control of Canada thistle and field bindweed.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

CORN

(FIELD, SEED*, POPCORN*, AND SILAGE CORN)

*Do not apply **Disha DMA** to seed corn or popcorn without first verifying with your local seed corn company (supplier) the **Disha DMA** selectively on your inbred line or variety of popcorn. This precaution will help avoid potential injury of sensitive varieties.

Disha DMA will control many annual broadleaf weeds or give growth suppression of many perennial broadleaf weeds commonly found in corn (refer to the **WEED LIST** table). For best performance, make application when weeds have emerged and are actively growing. Pre-emergence control of cocklebur, velvetleaf, and jimsonweed may be reduced if conditions such as low temperature or lack of soil moisture cause delayed or deep germination of weeds.

Agriculturally approved surfactants or sprayable fertilizers (½ to 1 gal. per acre of 28%, 30%, or 32% urea ammonium nitrate or 2.5 lbs. per acre spray grade ammonium sulfate**) may be added to the spray mixture to improve post-emergence weed control, particularly in dry growing conditions. **Not for use in California.

Precautions:

- Direct contact of **Disha DMA** with corn seed must be avoided.
- If corn seeds are less than 1 ½" below the surface, delay application until corn has emerged.
- Applications of **Disha DMA** to corn during periods of rapid growth may result in temporary leaning. Corn will usually become erect within 3 - 7 days.
- Cultivation should be delayed until after corn is growing normally to avoid breakage.
- For combination options or sequential treatments, refer to appropriate section.
- Several synthetic pyrethroid insecticides are labeled for tank mix applications of **Disha DMA**. Refer to their label for specific instructions.

Restrictions:

- Do not exceed a total of 1 ½ pts. (0.75 lb. a.e.) of **Disha DMA** per treated acre per crop year.
- Do not use adjuvants containing penetrants such as petroleum-based oils after crop emergence or crop injury may result.
- **Disha DMA** is not registered for use on sweet corn.
- Do not apply **Disha DMA** to seed corn or popcorn without first verifying with your local seed corn company (supplier) the selectivity of **Disha DMA** on your inbred line or variety of popcorn. This precaution will help avoid potential injury of sensitive varieties.
- Corn may be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity.
- Up to 2 applications of **Disha DMA** may be made during a growing season.
- Allow 2 weeks or more between applications of **Disha DMA**.

Pre-Plant/Pre-Emergence in No-Tillage Corn

Applications of **Disha DMA** may be made before, during, or after planting to emerged and actively growing broadleaf weeds. Apply **Disha DMA** at 1 pt. (0.5 lb. a.e.) per treated acre on medium- or fine-textured soils containing 2% or greater organic matter. Use ½ pt. (0.25 lb. a.e.) per treated acre on coarse-textured soils (sand, sandy loam, and loamy sand) or medium- and fine-textured soils with less than 2% organic matter.

When planting into a legume sod (e.g., alfalfa or clover), apply **Disha DMA** after 4 - 6" of regrowth has occurred.

Pre-Emergence in Conventional or Reduced Tillage Corn

Disha DMA may be applied after planting and prior to corn emergences. Application at 1 pt. (0.5 lb. a.e.) per treated acre may be made to medium- or fine-textured soils which contain 2% or greater organic matter. Do not apply to coarse-textured soils (sand, sandy loam, and loamy sand) until after crop emergence (refer to the **Early Post-Emergence** uses below).

Pre-emergence application of **Disha DMA** does not require mechanical incorporation to become active. A shallow mechanical incorporation is recommended if application is not followed by adequate rainfall or sprinkler irrigation. Avoid tillage equipment (e.g., drags, harrows) which concentrate treated soil over seed furrow.

Early Post-Emergence in All Tillage Systems (Spike through 8" tall corn)

Disha DMA at 1 pt. per treated acre may be applied during the period from corn emergence through the 5-leaf stage or 8" tall, whichever comes first. Reduce the rate to ½ pt. per treated acre if corn is growing on coarse-textured soils (sand, sandy loam, loamy sand). See **Late Post-Emergence** applications given below, if the 6th true-leaf is emerging from whorl or corn is greater than 8" tall.

Late Post-Emergence in All Tillage Systems (8 to 36" tall corn)

Application of **Disha DMA** at ½ pt. (0.25 lb. a.e.) per treated acre may be made from 8 - 36" tall corn or 15 days before tassel emergence, whichever comes first. For best performance, make applications when weeds are less than 3" tall.

Apply directed spray when corn leaves prevent proper spray coverage, sensitive crops are growing nearby, or tank mixing with 2,4-D.

Do not apply **Disha DMA** when soybeans are growing nearby if any of these conditions exist:

- Corn is more than 24" tall.
- Soybeans are more than 10" tall.
- Soybeans have begun to bloom.

Overlay (Sequential) Treatments

Disha DMA may be applied to ground previously treated with one or more of the following herbicides registered for use in corn:

acetochlor (Surpass [®] , Harness [®])	dimethenamid (Frontier [®])	Marksman [®]
alachlor (Lasso [®])	EPTC (Eradicane [®])	s-metolachlor (Dual [®] II Magnum)
atrazine	Guardsman	paraquat (Gramoxone [®])
Bicep II Magnum [®]	glyphosate (Roundup [®])	pendimethalin (Prowl [®])
Bullet [®]	halosulfuron (Permit [®])	simazine (Princep [®])
butylate (Sutan+ [®])	Lariat [®]	Surpass [®]
Clarity [®]		

Apply **Disha DMA** at ½ pt. (0.25 lb. a.e.) per treated acre to ground previously treated with full rates of Clarity or Marksman herbicides. Allow at least 2 weeks between applications.

Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Disha DMA may be tank mixed with one or more of, but not limited to, the following herbicides for control of grasses or additional broadleaf weeds in corn:

Disha DMA + (Plus)	Pre-Plant / Pre-E Mergence (No Tillage Corn)	Pre-Emergence (Conventional or Reduced Tillage Corn)	Early Post-Emergence (All Tillage Systems)	Late Post-Emergence (All Tillage Systems)	Additional Directions
Accent (nicosulfuron)	-	-	½ - 1.0 oz. a.e./A	½ - 1.0 oz. a.e./A To improve spray coverage of weeds and reduce risk of corn injury, use drop pipes to direct spray beneath corn leaves when corn is greater than 8" tall.	Application may be made to emerged weeds before corn is greater than 24" tall. Use non-ionic surfactant at 0.25% (v/v) with this tank mixture.
atrazine	1 ¼ - 2.0 lb. a.e./A	1 ¼ - 2.0 lb. a.e./A	1 ¼ - 2.0 lbs. a.e./A Crop oil concentrates may be used with this mixture if corn is 5" or less in height.	1 ¼ - 2.0 lbs. a.e./A Do not apply if corn is greater than 12" tall.	Application may be made before grasses are 1 ½" tall. Follow all State and Federal restrictions pertaining to atrazine applications.

Disha DMA + (Plus)	Pre-Plant / Pre-E Emergence (No Tillage Corn)	Pre-Emergence (Conventional or Reduced Tillage Corn)	Early Post-Emergence (All Tillage Systems)	Late Post-Emergence (All Tillage Systems)	Additional Directions
Beacon (primisulfuron)	-	-	0.31 - 0.62 oz. a.e./A	0.31 - 0.62 oz. a.e./A To improve spray coverage of weeds and reduce risk of corn injury, use drop pipes to direct spray beneath corn leaves when corn is greater than 8" tall.	Application may be made to emerged weeds when corn is 4 - 24" tall. Use non-ionic surfactant at 0.25% (v/v) with this tank mixture.
Dual II Magnum/ II Magnum SI (metolachlor)	0.8 - 1.6 lbs. a.e./A	0.8 - 1.6 lbs. a.e./A Use only on fine- or medium-textured soils with 2.5% or greater organic matter.	0.8 - 1.6 lbs. a.e./A	-	Application may be made before grasses reach the 2-leaf stage and before corn is greater than 3" tall.
Frontier (dimethenamid)	13.0 - 25.0 fl. oz./A	13.0 - 25.0 fl. oz./A Use only on fine- or medium-textured soils with 2.5% or greater organic matter.	13.0 - 25.0 fl. oz./A	-	Application may be made up to 8" tall corn. This treatment must be combined with a herbicide that provides post-emergence control of grass weeds if they are greater than 1" tall at the time of application.
Gramoxone (paraquat)	¼ - 1.0 lb. a.e./A	¼ - 1.0 lb. a.e./A	-	-	Application may be made to emerged weeds but prior to corn emergence.
Harness or Surpass (acetochlor)	1 ½ - 3.0 lbs. a.e./A	1 ½ - 3.0 lbs. a.e./A Use only on fine- or medium-textured soils with 2.5% or greater organic matter.	-	-	Application should be made prior to corn emergence.
Lasso (alachlor)	1 ½ - 4.0 lbs. a.e./A	1 ½ - 4.0 lbs. a.e./A Use only on textured soils with 2.5% organic matter.	1 ½ - 4.0 lbs. a.e./A	-	Application may be made before grasses reach the 2-leaf stage and before corn is greater than 3" tall. If micro-encapsulated forms of alachlor are used (Lasso MT, Partner), applications must be made prior to grass emergence.
Princep (simazine)	2.0 - 3.0 lbs. a.e./A	2.0 - 3.0 lbs. a.e./A	-	-	Application may be made prior to corn or weed emergence.
Prowl (pendimethalin)	-	¾ - 1 ½ lbs. a.e./A Use only on fine- or medium-textured soils with 2.5% or greater organic matter.	¾ - 1 ½ lbs. a.e./A	-	Application may be made immediately after planting but prior to weed emergence. Corn should not be beyond the 2-leaf stage of growth.
Roundup (glyphosate)	1.0 - 3.0 lbs. a.e./A	1.0 - 3.0 lbs. a.e./A	-	-	Application may be made to emerged weeds but prior to corn emergence.
2,4-D	¼ - ½ lb. a.e./A	¼ - ½ lb. a.e./A	Do not use at this application timing.	½ lb. a.e./A	Drop pipes are to be used when corn height is 8" or greater. Keeping the spray off the corn leaves and out of the whorl will reduce the likelihood of crop injury and improve spray coverage of the weed foliage.

COTTON

Pre-Plant Application

Apply **Disha DMA** as a broadcast or spot treatment to emerged and actively growing weeds at a rate of up to ½ pt. (0.25 lb. a.e.) per acre prior to planting cotton. Most effective control of weeds, apply **Disha DMA** when weeds are in the 2- to 4-leaf stage and rosettes are less than 2" across. Refer to the **WEED LIST** table for a list of weeds controlled or suppressed.

Note: Observe all precautions on this label, including references to crops growing under stress, drift, sensitive crops, and tank cleanout.

Restrictions - Cotton:

- Do not plant cotton for at least 21 days after application and after allowing for a minimum accumulation of 1" of rainfall or overhead irrigation.
- Do not apply to cotton west of the Rockies.
- Do not apply to cotton in geographic areas with average annual rainfall less than 25".
- If applying a Spring pre-plant treatment following application of a Fall pre-plant (post-harvest) treatment, the combination of both treatments may not exceed 4 pts. of this product (2 lbs. a.e.) per year.

Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

For control of grasses or additional broadleaf weeds, **Disha DMA** may be tank mixed with Caparol, Gramoxone Extra, and Roundup Ultra RT herbicides.

SORGHUM (MILO)

Disha DMA when applied at the specified rate for sorghum, will control many actively growing annual broadleaf weeds and will reduce competition from established perennial broadleaf weeds as well as control their seedlings (refer to the **WEED LIST** table).

Note: Observe all precautions. Read and follow mixing, application, and cleaning instructions.

If sorghum is grown for pasture or hay, refer to the **PASTURE, HAY, RANGELAND, AND FARMSTEAD (NON-CROPLAND)** use section of the label.

Restrictions:

- Applications of **Disha DMA** to sorghum during periods of rapid growth may result in temporary leaning of plants or rolling of leaves. These effects are usually outgrown within 10 - 14 days.
- Do not graze or feed treated sorghum forage or silage prior to mature grain stage.
- Do not apply **Disha DMA** to sorghum grown for seed production.
- Make no more than 1 application per growing season.

Pre-Plant Applications

½ pt. (0.25 lb. a.e.) of **Disha DMA** may be applied to emerged and actively growing weeds at least 15 days prior to planting.

Post-Emergence Applications

½ pt. (0.25 lb. a.e.) of **Disha DMA** must be made after sorghum is in the spike stage (all sorghum emerged), but before sorghum is 15" tall. For best performance, make applications when sorghum is in the 3- to 5-leaf stage and weeds are small (less than 3" tall). Use drop pipes (drop nozzles) if sorghum is taller than 8". Keeping the spray off the sorghum leaves and out of the whorl will reduce the likelihood of crop injury and improve spray coverage of weed foliage.

Pre-Harvest in Texas and Oklahoma Only

½ pt. (¼ lb. a.e.) of **Disha DMA** may be applied for weed suppression any time after the sorghum has reached the soft dough stage. An agriculturally approved surfactant may be used to improve performance. For aerial applications, use at least 2 gals. of water-based carrier per treated acre. Delay harvest until 30 days after treatment.

Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Disha DMA plus Atrazine

For improved control of emerged, actively growing broadleaf weeds including triazine-resistant species and added suppression of perennial broadleaf weeds, tank mix ½ pt. (0.25 lb. a.e.) **Disha DMA** with labeled use rates of atrazine per treated acre. For control of grasses (less than 1.5" tall), tank mix ½ pt. (0.25 lb. a.e.) **Disha DMA** with labeled use rate of atrazine per treated acre. For best performance and minimal crop injury, make application when sorghum is 3 - 8" tall and when broadleaf weeds are small (less than 6" tall). Application of atrazine must be made before sorghum is beyond 12" tall. The atrazine rate will depend upon soil texture and length of residual weed control desired. Follow all State and Federal restrictions pertaining to atrazine applications.

Disha DMA plus Buctril® (bromoxynil)

For improved control of broadleaf weeds, tank mix ½ pt. (0.25 lb. a.e.) **Disha DMA** with labeled use rates of Buctril per treated acre. Make application at 4-leaf to 15" tall sorghum. Use drop nozzles to direct spray beneath sorghum leaves when sorghum is greater than 8" tall.

Overlay (Sequential) Treatments

Disha DMA may be applied to ground previously treated with one or more of the following herbicides:

Herbicide	Maximum Rate per Treated Acre (Lb. A.E.)
atrazine ¹	2.5
s-metolachlor (Dual II Magnum®) (Concep® treated seed)	1.33

¹Maximum use rate for atrazine is determined by soil type, tillage practices used, surface residue, and State or local restrictions. Follow the more restrictive requirements when determining the maximum use rate for atrazine.

SMALL GRAINS (WHEAT, BARLEY, AND OATS)

Not Underseeded to Legumes.

Application of **Disha DMA** may be made before, during or after planting of small grains. For best performance, make applications when weeds are in the 2- to 3-leaf stage and rosettes are less than 2" across. Application of **Disha DMA** to small grains during periods of rapid growth may result in crop leaning. This condition is temporary and will not reduce crop yields.

Use **Disha DMA** at 2 - 4 fl. oz. (0.063 to 0.125 lb. a.e.) per treated acre in wheat, fall seeded barley, and oats, and at 2 - 3 fl. oz. (0.063 to 0.094 lb. a.e.) per treated acre in spring seeded barley. Use the higher level of listed rate ranges when treating difficult to control weeds such as kochia, Russian thistle, and prickly lettuce or dense vegetative growth.

Disha DMA used in a tank mix with other herbicides offers the best spectrum of weed control and herbicide tolerant or resistant weed management. Refer to specific crop for **Disha DMA** rate and application timing.

Note: Observe all precautions. Read and follow cleaning, mixing, application, and cleaning instructions.

Restrictions:

- Pre-harvest interval (PHI) restriction for grain is 7 days.
- **Disha DMA** contains 0.5 lb. a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 lb. of a.e. per acre per application.
- If small grains are used for pasture hay, the following restrictions apply:
 - Animals cannot be removed from treated area for slaughter prior to 30 days after last application.
 - There is no waiting period between treatment and grazing for non-lactating dairy animals.
 - Treated areas may not be grazed by lactating dairy animals before 7 days after treatment.
 - Do not harvest hay from treated areas before 37 days after treatment.

Weeds Controlled

Disha DMA or combinations with listed tank mix partners, will provide control or suppression of annual broadleaf weeds listed below. For improved control of listed weeds, it is recommended that **Disha DMA** be applied in a tank mix with other herbicides. Refer to specific crop for tank mix options.

Alkanet ¹	Henbit	Pineappleweed ¹
Bedstraw, Catchweed ¹	Jacob's Ladder ¹	Plantain, Broadleaf ²
Bindweed, Field ²	Knawel (German Moss)	Poppy, Red Horned ¹
Buckwheat, Tartary	Knotweed, Prostrate	Puncturevine ¹
Buckwheat, Wild	Kochia	Puslane, Common ¹
Carpetweed ¹	Ladysthumb	Radish, Wild ¹
Chamomile, Corn	Lambsquarters, Common	Ragweed, Common ¹
Chervil, Bar	Lettuce, Miners ¹	Ragweed, Giant (Buffaloweed) ¹
Chickweed, Common ¹	Lettuce, Prickly	Rocket, London ¹
Cockle, Corn	Mallow, Common	Rocket, Yellow ¹
Cockle, Cow	Mayweed, Chamomile (Dogfennel)	Salsify (Goatsbeard) ¹
Cocklebur, Common	Mustard, Blue (Purple)	Shepherd's Purse ¹
Cornflower (Bachelorbutton) ¹	Mustard, Tansy	Smartweed, Green
Dandelion, Common ²	Mustard, Treacle ¹	Smartweed, Pennsylvania
Dock, Curly ²	Mustard, Tumble (Jim Hill) ¹	Sorrel, Red (Sheep Sorrell) ¹
Dragonhead, American ¹	Mustard, Wild ¹	Sowthistle, Annual
Evening Primrose, Cutleaf ¹	Nightshade, Black	Starthistle, Yellow ¹
Falseflax, Smallseeded ¹	Nightshade, Cutleaf ¹	Sunflower, Common (Wild)
Fiddleneck (Tarweed) ¹	Nightshade, Silverleaf ² (White Horsenettle)	Thistle, Canada ²
Flixweed ¹	Pennycress, Field (Fanweed, Frenchweed, Stinkweed)	Thistle, Russian
Fumitory ¹	Pepperweed, Peppergrass ¹	Velvetleaf
Gromwell, Corn ¹	Pigweed, Redroot (Carelessweed)	Vetch ¹
Groundsel, Common ¹	Pigweed, Rough	Yarrow, Common ²
Hempnettle ¹	Pigweed, Tumble	

¹These weeds will be controlled with **Disha DMA** tank mixtures. Refer to tank mix label for specific weeds controlled.

²**Disha DMA** tank mixes will provide suppression of established broadleaf weeds and control of their seedlings.

Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

For applications prior to the emergence of weeds or when sulfonylurea-resistant weeds are present or suspected, use a minimum of 3 fl. oz. (0.094 lb. a.e.) per treated acre of **Disha DMA** with a tank mix herbicide. Non-sulfonylurea herbicides such as 2,4-D or MCPA tank mixed with **Disha DMA** will offer more consistent control of sulfonylurea-resistant weeds.

When tank mixing with sulfonylurea herbicides, such as Ally[®], Amber[®], Express[®], and Harmony[®] Extra use an agriculturally approved surfactant of at least 80% active ingredient at the rate of 1 - 4 pts./100 gals. of spray or not more than 0.25 - 0.5% by volume. Use the highest rate of surfactant when using the lower rate ranges of the tank mix and/or when treating more mature and difficult to control weeds or dense vegetative growth.

FALL AND SPRING SEEDED WHEAT

DISHA DMA MUST BE APPLIED TO FALL SEEDED WHEAT PRIOR TO THE JOINTING STAGE. APPLICATIONS TO SPRING SEEDED WHEAT MUST BE MADE BEFORE WHEAT EXCEEDS THE 5-LEAF STAGE. Early developing wheat varieties such as TAM107, MADISON, or WAKEFIELD must receive application between early tillering and the jointing stage. Care should be taken in staging these varieties to be certain that the application occurs prior to the jointing stage.

Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Disha DMA may be tank mixed with one or more, but not limited to, the following herbicides.

Broadcast Rate per Treated Acre – Apply 2.0 - 4.0 fl. oz. (0.063 to 0.125 lb. a.e) **Disha DMA** with:

Product	Active Ingredient	Formulation	Amount of Product per Acre
2,4-D amine or ester	2,4-D	4.0 lbs./gal.	8.0 - 12.0 fl. oz. (0.25 - 0.375 lb. a.e./A)*
MCPA amine or ester	MCPA	4.0 lbs./gal.	8.0 - 12.0 fl. oz. (0.25 - 0.375 lb. a.e./A)*
Ally®	metsulfuron-methyl	60% DF	⅓ oz.
Amber®	triasulfuron	75% DF	0.28 oz.
Express®	thifensulfuron + tribenuron-methyl	75% DF	⅓ oz.
Harmony® Extra	thifensulfuron + tribenuron-methyl	75% DF	⅓ oz.
Buctril®	bromoxynil ¹	2.0 lbs./gal.	1.0 - 1.5 pts.
Bronate®	bromoxynil + MCPA	4.0 lbs./gal.	1.0 - 2.0 pts.
Curtail®	clopyralid + 2,4-D	2.38 lbs./gal.	2.0 - 2 ⅓ pts.
Stinger	clopyralid	3.0 lbs./gal.	¼ - ½ pts.
Karmex® ²	diuron ¹	80% DF	½ - 1.5 lbs.
Sencor® ²	metribuzin ¹	75% DF	1.0 - 10.0 oz.

*When using formulations other than 4.0 lbs./gal., use pounds active/acre listed.

¹Herbicides with the same active ingredient and/or different formulations may be used.

²Tank mixtures for fall seeded wheat only.

Special Use Tank Mixes For Spring And Fall Seeded Wheat (See Footnotes For Applicable Uses.)

Broadcast Rate per Treated Acre – Apply 3.0 - 4.0* fl. oz. (0.094 to 0.125 lb. a.e) of **Disha DMA** with:

Product ¹	Active Ingredient	Amount of Formulation	Product per Acre
2,4-D or MCPA amine	2,4-D or MCPA	4.0 lbs./gal.	1.0 - 2.0 pts. ² (0.5 - 1.0 lb. a.e./A) ³
2,4-D or MCPA ester	2,4-D or MCPA	4.0 lbs./gal.	1.0 - 1.5 pts. ² (0.5 - 0.75 lb. a.e./A) ³
Ally	metsulfuron-methyl	60% DF	⅒ - ⅓ oz.
Amber	triasulfuron	75% DF	0.14 - 0.28 oz.
Express	thifensulfuron + tribenuron-methyl	75% DF	⅓ - ⅓ oz.
Harmony Extra	thifensulfuron + tribenuron-methyl	75% DF	⅓ - ⅓ oz.
Ally + 2,4-D amine or ester ⁴	metsulfuron-methyl + 2,4-D	60% DF + 4.0 lbs./gal.	⅒ - ⅓ oz. + 8.0 fl. oz.
Amber + 2,4-D amine or ester ⁴	triasulfuron + 2,4-D	75% DF + 4.0 lbs./gal.	0.14 oz. - 0.28 oz. + 8.0 fl. oz.
Express + 2,4-D amine or ester ⁴	(thifensulfuron + tribenuron-methyl) + 2,4-D	75% DF + 4.0 lbs./gal.	⅓ - ⅓ oz. + 8.0 fl. oz.
Harmony Extra + 2,4-D amine or ester ⁴	(thifensulfuron + tribenuron-methyl) + 2,4-D	75% DF + 4.0 lbs./gal.	⅓ - ⅓ oz. + 8.0 fl. oz.
Roundup ⁵	glyphosate	3.0 lbs./gal.	12.0 - 16.0 fl. oz.

Disha DMA may be used at 6.0 fl. oz. (0.22 lb. a.e.) on fall seeded wheat in Western Oregon as a spring application only. In CO, KS, NM, OK, and TX up to 8 fl. oz. of **Disha DMA** may be applied on fall seeded wheat after it exceeds the 3-leaf stage for suppression of perennial weeds, such as field bindweed. Applications may be made in the fall following a frost but before a killing freeze. **Disha DMA** may be tank mixed with 2,4-D amine (see label for use rates) after wheat begins to tiller. Periods of extended stress such as cold and wet weather may enhance the possibility of crop injury. For fall applications only, do not use if the potential for crop injury is not acceptable.

¹Do not use low rates of sulfonyleurea herbicides, such as Ally, Amber, Express, and Harmony Extra on more mature weeds and/or on dense vegetative growth.

²**Note:** For use on Fall Seeded Wheat only. Do not use unless potential crop injury will be acceptable.

³When using formulations other than 4.0 lbs./gal., use pounds active/acre listed.

⁴Use for improved control of Russian thistle, flaxweed, gromwell, mayweed, and fiddleneck.

⁵**Disha DMA** may be applied at 2.0 fl. oz. with Roundup as a pre-plant application to small grains with no waiting period prior to planting. Add 0.5% by volume of an agriculturally approved non-ionic surfactant.

FALL SEEDED BARLEY

DISHA DMA MUST BE APPLIED TO FALL SEEDED BARLEY PRIOR TO THE JOINTING STAGE.

Note: For spring barley varieties that are seeded during the winter months or later, follow the rates and timings given for spring seeded barley.

Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Disha DMA may be tank mixed with one or more of the following herbicides.

Broadcast Rate per Treated Acre – Apply 2.0 - 4.0 fl. oz. (0.063 to 0.125 lb. a.e) **Disha DMA** with:

Product ¹	Active Ingredient	Formulation	Amount of Product per Acre
2,4-D amine or ester	2,4-D	4.0 lbs./gal.	Refer to product label
MCPA amine or ester	MCPA	4.0 lbs./gal.	Refer to product label
Ally®	metsulfuron-methyl	60% DF	Refer to product label
Amber®	triasulfuron	75% DF	Refer to product label
Express®	thifensulfuron + tribenuron-methyl	75% DF	Refer to product label
Harmony® Extra	thifensulfuron + tribenuron-methyl	75% DF	Refer to product label
Sencor®	metribuzin ²	75% DF	Refer to product label
Buctril®	bromoxynil	2.0 lbs./gal.	Refer to product label
Bronate®	bromoxynil + MCPA	4.0 lbs./gal.	Refer to product label

¹Do not use low rates of sulfonylureas (Ally, Amber, Express, and Harmony Extra) on more mature weeds and/or on dense vegetative growth.

²Herbicides with the same active ingredient and/or different formulations may be used.

SPRING SEEDED BARLEY

DISHA DMA MUST BE APPLIED BEFORE SPRING SEEDED BARLEY EXCEEDS THE 4-LEAF STAGE.

Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Disha DMA may be tank mixed with one or more of the following herbicides.

Broadcast Rate per Treated Acre – Apply 2.0 - 3.0 fl. oz. (0.063 to 0.094 lb. a.e) **Disha DMA** with:

Product ¹	Active Ingredient	Formulation	Amount of Product per Acre
MCPA amine or ester	MCPA	4.0 lbs./gal.	Refer to product label
Ally®	metsulfuron-methyl	60% DF	Refer to product label
Amber®	triasulfuron	75% DF	Refer to product label
Express®	thifensulfuron + tribenuron-methyl	75% DF	Refer to product label
Harmony® Extra	thifensulfuron + tribenuron-methyl	75% DF	Refer to product label
Sencor®	metribuzin ²	75% DF	Refer to product label
Buctril®	bromoxynil	2.0 lbs./gal.	Refer to product label
Bronate®	bromoxynil + MCPA	4.0 lbs./gal.	Refer to product label

¹Do not use low rates of sulfonylureas (Ally, Amber, Express, and Harmony Extra) on more mature weeds and/or on dense vegetative growth.

²Herbicides with the same active ingredient and/or different formulations may be used.

FALL AND SPRING SEEDED OATS

DISHA DMA MUST BE APPLIED BEFORE SPRING SEEDED OATS EXCEED THE 5-LEAF STAGE. APPLICATIONS TO FALL SEEDED OATS MUST BE MADE PRIOR TO THE JOINTING STAGE.

Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Disha DMA may be tank mixed with one or more of the following herbicides.

Broadcast Rate per Treated Acre – Apply 2.0 - 4.0 fl. oz. (0.063 to 0.125 lb. a.e) **Disha DMA** with:

Product	Active Ingredient	Formulation	Amount of Product per Acre
MCPA amine or ester	MCPA	4.0 lbs./gal.	Refer to product label

SUGARCANE

Disha DMA when applied at specified rates, will control many annual, biennial, and perennial broadleaf weeds commonly found in sugarcane (refer to the **WEED LIST** table).

Application of **Disha DMA** may be made any time after weeds have emerged and are actively growing but before the close-in stage of sugarcane.

When possible, direct the spray beneath the sugarcane canopy in order to minimize the likelihood of crop injury. The use of directed sprays will also aid in maximizing spray coverage of weed foliage.

Consult your local or State authorities for possible application restrictions, especially concerning aerial applications and advice concerning special local use situations.

Note: Observe all precautions. Read and follow mixing, application, and cleaning instructions.

Restrictions:

- Do not apply within 87 days of harvest.
- Retreatment may be made as needed, however, do not exceed a total of 2 applications of **Disha DMA** per year.

Application rates and timings of **Disha DMA** are given below. Use the higher level of listed rate ranges when treating dense vegetative growth.

Weed Stage & Type	Broadcast Rate per Treated Acre	
	Amount of Product	Equivalent Lb. A.E.
Annual		
Small, actively growing	½ - 1.0 pt.	¼ - ½
Established weed growth	1.0 - 1 ½ pts.	½ - ¾
Biennial	1.0 - 2.0 pts.	½ - 1.0
Perennial	2.0 pts.	1.0*

*Application made over the top of actively growing sugarcane may result in crop injury.

Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Disha DMA may be tank mixed with one or more of, not limited to, the following herbicides for control of grasses or additional broadleaf weeds:

Herbicide	Rates per Treated Acre
ametryn (Evik®)	Refer to product label
asulam (Asulox®)	Refer to product label
atrazine	Refer to product label
2,4-D*	Refer to product label

*Application of **Disha DMA** plus 2,4-D tank mix at the higher level rate ranges may result in crop injury.

PASTURE, HAY, RANGELAND, AND FARMSTEAD (NON-CROPLAND)

Disha DMA, when applied at specified rates, will give control of many annual, biennial, and perennial broadleaf weeds, and many woody brush and vine species commonly found in pasture, hay, rangeland, and general farmstead (non-cropland) areas (refer to the **WEED LIST** table). Perennial weeds noted with a superscript 1 in the **WEED LIST** table may be controlled with lower rates of either **Disha DMA** or **Disha DMA** plus 2,4-D.

Disha DMA may also be applied to non-cropland areas for the control of broadleaf weeds in Noxious Weed Control Programs, Districts or Areas including broadcast or spot treatment of roadsides and highways, utilities, railroad, and pipeline rights-of-way. Noxious weeds must be recognized at the State level but programs may be administered at State, County, or other levels.

Disha DMA uses described in this section also pertain to small grains (such as barley, forage sorghum, oats, rye, sudangrass or wheat) grown for pasture use only.

Note: Observe all precautions. Read and follow mixing, application, and cleaning instructions.

Restrictions:

- The pre-harvest interval (PHI) for grass forage is 0 days.
- The pre-harvest interval (PHI) for grass hay is 7 days.
- Retreatments may be made as needed; however, do not exceed a total of 2 applications of **Disha DMA** per year.
- Newly Seeded Areas may be severely injured if rates of **Disha DMA** are greater than 1 pt./A.
- Established Grass Crops growing under stress can exhibit various injury symptoms that may be more pronounced if herbicides are applied.
- Bentgrass, carpetgrass, buffalograss, and St. Augustinegrass may be injured at rates exceeding 1 pt. **Disha DMA** (½ lb. a.e.) per treated acre. Usually colonial bentgrasses are more tolerant than creeping types. Velvetgrasses are most easily injured. Treatments will kill or injure alfalfa, clovers, lespedeza, wild winter peas, vetch, and other legumes.
- Animals cannot be removed from treated area for slaughter prior to 30 days after last application. There is no waiting period between treatment and grazing for non-lactating animals.

Timing Restrictions For Lactating Dairy Animals Following Treatment		
Disha DMA Rate per Treated Acre	Days Before Grazing (Days)	Days Before Hay Harvest (Days)
Up to 1.0 pt. (½ lb. a.e.)	7	37
Up to 2.0 pts. (1 lb. a.e.)	21	51

Application rates and timing of **Disha DMA** are given in the below table. Use the higher level of listed rate ranges when treating dense or tall vegetative growth.

Weed Stage & Type	Broadcast Rate per Treated Acre	
	Amount of Product	Equivalent Lb. A.E.
Annuals		
Small, actively growing	½ - 1.0 pt.	¼ - ½
Established weed growth	1.0 - 1 ½ pts.	½ - ¾
Biennials¹ (Rosette diameter)		
Less than 3"	½ - 1.0 pt.	¼ - ½
3" or more	1.0 - 2.0 pts.	½ - 1.0
Bolting	2.0 pts.	1.0

¹For best performance, make application when Biennial weeds are in the rosette stage.

(continued)

Weed Stage & Type	Broadcast Rate per Treated Acre	
	Amount of Product	Equivalent Lb. A.E.
Perennials		
Suppression or top growth control	½ - 1.0 qt.	½ - 1.0
Noted perennials (Perennial weeds noted with a superscript 1 in the WEED LIST table.)	1.0 qt.	1.0
Other Perennials	1.0 qt.	1.0
Woody Brush & Vines		
Top growth suppression	½ - 1.0 qt.	½ - 1.0
Top growth control ²	1.0 qt.	1.0

²Species noted in **WEED LIST** table will require tank mixtures for adequate control.

Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Disha DMA can be applied using water, oil-in water emulsions (including invert systems), or sprayable fluid fertilizer as a carrier. A compatibility test (refer to the **Compatibility Test** section) should be made prior to tank mixing.

To prepare oil-in water emulsions, half-fill spray tank with water, then add the appropriate amount of emulsifier. With continuous agitation, slowly add the herbicide and then the oil (such as diesel oil or fuel oil) or a premix of oil plus additional emulsifier to spray tank. Complete filling of spray tank with water. Maintain vigorous agitation during spray operation to prevent oil and water from forming separate layers.

Disha DMA may be applied broadcast using either ground or aerial application equipment. When using ground equipment, apply 3 - 600 gallons of diluted spray per treated acre. Volume of spray applied will depend on the height, density, and type of weeds or brush being treated and on the type of equipment being used. When using aerial equipment, apply 1 - 40 gallons of diluted spray per treated acre in a water-based carrier.

Disha DMA may be applied to individual clumps or small areas (spot treatment) of undesirable vegetation using handgun or similar types of application equipment. Apply diluted sprays to allow complete wetting (up to runoff) of foliage and stems.

Herbicide adjuvants or other spray additives (emulsifiers, surfactants, wetting agents, drift control agents, or penetrants) may be used for wetting, penetration, or drift control. Spray additives must be agriculturally approved when used in pasture applications. If spray additives are used, read and follow all use recommendations and precautions on product label.

Disha DMA may be tank mixed with one or more, but not limited to, of the following herbicides for control of grasses, additional broadleaf weeds, and woody brush and vines.

Herbicide	Rates per Treated Acre
Pasture, hay, rangeland, and general farmstead (non-cropland) use:	
glyphosate (Roundup®)	Refer to product label
metsulfuron-methyl (Ally)	Refer to product label
paraquat (Gramoxone)	Refer to product label
picloram (Tordon®)	Refer to product label
triclopyr (Garlon®)	Refer to product label
2,4-D	Refer to product label

Due to the variations that may occur in formulated products and specific use ingredients (e.g., water supplies), a compatibility test (refer to the **Compatibility Test** section) is recommended prior to actual tank mixing.

CUT SURFACE TREE TREATMENTS

Disha DMA may be applied as a cut surface treatment for control of unwanted trees and prevention of sprouts of cut trees. Use a mix of 1 part **Disha DMA** with 1 - 3 parts water. Use the lower dilution when treating difficult-to-control species.

- **Frill or Girdle Treatments:** Make a continuous cut or a series of overlapping cuts using an axe to girdle tree trunk. Spray or paint cut surface with the **Disha DMA**/water mix.
- **Stump Treatments:** Spray or paint freshly cut surface with the water mix. The area adjacent to the bark should be thoroughly wet.

Note: For more rapid foliar effects, 2,4-D may be added to the **Disha DMA**/water mix.

DORMANT APPLICATIONS FOR CONTROL OF MULTIFLORA ROSE

Disha DMA can be applied when plants are dormant as an undiluted Spot Concentrate directly to the soil or as a Lo-Oil basal bark treatment using an oil-water emulsion solution.

- **Spot Concentrate Treatments:** Apply **Disha DMA** directly to the soil as close as possible to the root crown but within 6 - 8" of the crown. On sloping terrain, make application to the uphill side of the crown. Do not make application when snow or water prevents applying **Disha DMA** directly to the soil. The use rate of **Disha DMA** is dependent on the canopy diameter of the multiflora rose. Examples: Use **Disha DMA** at ¼, 1.0, or 2 ¼ fl. oz. of product respectively, for 5, 10, or 15 feet canopy diameters. Do not exceed a total of 2 qts. **Disha DMA** per acre per year.
- **Lo-Oil Basal Bark Treatments:** Apply **Disha DMA** to the basal stem region from the ground line up to a height of 12 - 18". Spray until runoff, with special emphasis on covering the root crown. For best results, make application when plants are dormant. Do not make application after bud break or when plants are showing signs of active growth. Do not make application when snow or water prevents applying **Disha DMA** to the ground line.

Refer to **Tank Mixtures** in this section for method of preparing oil-in-water emulsion.

Example for making approximately 2 gallons of a LO-OIL spray solution mixture:

1. Combine 1 ½ gals. of water plus 1 oz. emulsifier plus 1 pt. (0.5 lb. a.e.) **Disha DMA** plus 2 ½ pts. of No. 2 diesel fuel.
2. Adjust amounts of materials used proportionately to the amount of final spray solution desired. Do not exceed 8 gals. of final spray solution applied per acre per year.

CONSERVATION RESERVE PROGRAM (CRP) ACRES

Disha DMA is recommended for use on both newly seeded and established grasses grown in Conservation Reserve or Federal Set-A-Side Programs.

Note: Observe all precautions. Read and follow mixing, application, and cleaning instructions.

Restrictions:

- **Disha DMA** treatment will cause injury or may kill alfalfa, clovers, lespedeza, wild winter peas, vetch, and other legumes.
- Agriculturally approved surfactants may be added to the spray mixture to improve post-emergence weed control, particularly in dry growing conditions.
- Do not use adjuvants containing penetrants such as petroleum-based oils after grass emergence on newly seeded grasses.

NEWLY SEEDED AREAS

Disha DMA may be applied either pre-plant or post-emergence to newly seeded grasses or small grains such as barley, oats, rye, sudangrass, wheat or other grain species grown as a cover crop. Post-emergence applications may be made after seedling grasses exceed the 3-leaf stage. Rates of **Disha DMA** greater than 1 pt. (0.5 lb. a.e.) per treated acre may severely injure newly seeded grasses. Pre-plant applications may injure to new seedlings may occur if intervals between application and grass planting is less than 45 days per pint of **Disha DMA** per treated acre west of the Mississippi River or 20 days per pt. east of the Mississippi River.

ESTABLISHED GRASS STANDS

Disha DMA, when applied at specified rates, will control many annual and biennial weeds and provide control or suppression of many perennial weeds (refer to the **WEED LIST** table).

Established grass stands are perennial grasses planted one or more seasons prior to treatment. Certain species: bentgrass, carpetgrass, smooth brome, buffalograss or St. Augustinegrass may be injured when treated with **Disha DMA** at rates exceeding 1 pt. per treated acre.

Application rates and timing of **Disha DMA** treatment are given below. Use the higher rate of the rate range when vegetation is either dense or tall, or when weeds are growing under stressed conditions such as drought or cool temperature.

Weed Type ¹ & Stage	Broadcast Rate per Treated Acre	
	Amount of Product	Equivalent Lb. A.E.
Annuals		
Small, actively growing	¼ - 1.0 pt.	⅙ - ½
Established weed growth	1.0 pt.	½
Biennials² (Rosette diameter)		
Less than 3"	½ - 1.0 pt.	¼ - ½
3" or more	1.0 - 2.0 pts.	½ - 1.0
Bolting	2.0 pts.	1.0
Perennials²		
Suppression/Control	2.0 pts.	1.0

¹For best results, treat biennial weeds with **Disha DMA** when they are in the rosette stage of growth. Retreatment may be made as needed; however, do not exceed a total of 2 applications of **Disha DMA** per treated acre during a growing season.

²Biennial and perennial weeds will require follow-up (sequential) treatments for seedling control and escapes.

Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

To control grasses and additional broadleaf weeds, **Disha DMA** may be tank mixed with other herbicides registered for use in Conservation Reserve Programs such as 2,4-D, glyphosate (Roundup), paraquat (Gramoxone), metsulfuron (Ally) and others.

TURF AND LAWNS FOR USE IN FARMSTEAD (NON-CROPLAND) AND SOD FARMS

Disha DMA, when applied at specified rates, will give control of many annual, biennial, and perennial broadleaf weeds (noted with a superscript 1 in the **WEED LIST** table) commonly found in turf. **Disha DMA** will also give growth suppression of many other listed perennial broadleaf weeds and woody brush and vine species (refer to **WEED LIST** table).

Apply 30 - 200 gals. of diluted spray per treated acre (3 qts. - 4 ¼ gals. per 1,000 sq. ft.), depending on density or height of weeds treated and on the type of equipment used.

For best performance, apply when weeds are emerged and actively growing.

Retreatment may be made as needed, however, do not exceed a total of 2 applications of **Disha DMA** per year.

Note: Observe all precautions. Read and follow mixing, application, and cleaning instructions.

Restrictions:

- Do not exceed a total of 2 pts. (1 lb. a.e.) **Disha DMA** per treated acre during the growing season.
- To avoid injury to newly seeded grasses, application of **Disha DMA** should be delayed until after the second mowing. Furthermore, application rates in excess of 1 pt. (½ lb. a.e.) per treated acre may cause noticeable stunting or discoloration of sensitive grass species such as bentgrass, carpetgrass, buffalograss, and St. Augustinegrass.
- In areas where roots of sensitive plants extend, do not apply in excess of ¼ pt. (0.125 lb. a.e.) of **Disha DMA** per treated acre on coarse-textured (sandy-type) soils, or in excess of ½ pt. (0.25 lb. a.e.) per treated acre on fine-textured (clay-type) soils. Do not make repeat applications in these areas for 30 days and until previous applications of **Disha DMA** have been activated in the soil by rain or irrigation.

Use the higher level of listed rate ranges when treating dense vegetative growth.

Weed Stage & Type	Disha DMA		
	Pints per Treated Acre	Lb. A.E. per Treated Acre	Teaspoons per 1,000 Sq. Ft.
Annuals			
Small, actively growing	½ - 1.0	¼ - ½	1.0 - 2 ¼
Established weed growth	1.0 - 1 ½	½ - ¾	2 ¼ - 3 ¼
Biennials (Rosette diameter)			
Less than 3"	½ - 1.0	¼ - ½	1.0 - 2 ¼
3" or more	1.0 - 2.0	½ - 1.0	2 ¼ - 4 ½
Perennials and Woody Brush and Vines	1.0 - 2.0	½ - 1.0	2 ¼ - 4 ½

Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Tank mix treatments of **Disha DMA** may be made with 2,4-D, MCPA, MCPP, or bromoxynil for control of additional weeds listed on the tank mix product label.

Apply 3.2 to 8 fl. oz. (0.1 - 0.25 lb. a.e.) of **Disha DMA** per treated acre with labeled use rates of 2,4-D, MCPA, or MCPP, or with labeled use rate of bromoxynil. Use the higher level of the listed rate ranges when treating established weeds. Repeat treatments may be made as needed; however, do not exceed 2 pts. (1 lb. a.e.) of **Disha DMA** per treated acre during the growing season.

GRASS SEED CROPS (BERMUDA GRASS, BLUEGRASS, FESCUE, AND RYEGRASS)

Apply ½ - 1 pt. (0.25 to 0.5 lb. a.e.) of **Disha DMA** per treated acre on seedling grass after the crop reaches the 3- to 5-leaf stage. Apply up to 2 pts. (1 lb. a.e.) of **Disha DMA** on well-established perennial grass. For best performance, make applications when weeds are in the 2- to 4-leaf stage and rosettes are less than 2" across. Use the higher level of listed rate ranges when treating more mature weeds or dense vegetative growth.

Refer to the **PASTURE, HAY, RANGELAND, AND FARMSTEAD (NON-CROPLAND)** section for possible grazing and feeding restrictions.

Note: Observe all precautions. Read and follow mixing, application, and cleaning instructions.

Restrictions:

- Do not use on bentgrass unless possible crop injury can be tolerated.
- Do not apply after the grass seed crop begins to joint.

Weeds Controlled

Disha DMA will provide control or suppression of annual broadleaf weeds listed below. For improved control of listed weeds plus additional weeds, it is recommended that **Disha DMA** be applied in a tank mix with other herbicides.

Alfalfa ¹	Clover	Ladysthumb
Bedstraw, Catchweed	Cockle, White	Lambsquarters, Common
Bindweed, Field	Dock, Broadleaf	Lettuce, Prickly
Buttercup, Corn	Dock, Curly	Mayweed (Dogfennel)
Buttercup, Creeping	Hemlock, Poison	Ragwort, Tansy
Buttercup, Western Field	Knapweed, Russian ¹	Sorrel, Red (Sheep Sorrel)
Catchfly, Nightflowering	Knawel	Sowthistle, Annual
Chamomile, Corn	Knotweed, Prostrate	Starwort, Little
Chickweed, Common	Kochia	Thistle, Canada ¹
Chickweed, Mouseear		
¹ Top growth only.		

Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

For control of grasses or additional broadleaf weeds, **Disha DMA** may be tank mixed with all broadleaf herbicides registered for use in Grass Seed Production.

Broadcast Rate per Treated Acre – Apply ½ - 2.0 pts. (0.25 to 1 lb. a.e.) **Disha DMA** with:

Product	Active Ingredient	Formulation	Amount of Product per Acre
2,4-D amine or ester	2,4-D	4.0 lbs./gal.	Refer to product label
MCPA amine	MCPA	4.0 lbs./gal.	Refer to product label
Buctril®	bromoxynil	2.0 lbs./gal.	Refer to product label
Curtail®	clopyralid + 2,4-D	2.38 lbs./gal.	Refer to product label
Karmex®	diuron	80% DF	Refer to product label
Stinger®	clopyralid	3.0 lbs./gal.	Refer to product label

ANNUAL GRASS CONTROL

For suppression of annual grass weeds such as:
Downy Brome (Cheatgrass), Rippgut Brome, Rattail Fescue, and Windgrass

Apply 2 pts. (1 lb. a.e.) of **Disha DMA** per treated acre in the fall or late summer after harvest and burning of established grass seed crops. Applications should be made immediately following the first irrigation when the soil is moist and before weeds have more than 2 leaves.

**PRE-PLANT DIRECTIONS
(POST HARVEST/ FALLOW/CROP STUBBLE/SET-A-SIDE)
FOR BROADLEAF WEED CONTROL BEFORE WHEAT, CORN, SORGHUM, SOYBEANS**

Apply **Disha DMA** as a broadcast or spot treatment to emerged and actively growing weeds after crop harvest (post-harvest) and before a killing frost or in the fallow cropland or crop stubble the following spring or summer. Agriculturally approved spray additives, such as surfactants or oils, may be used to enhance spray coverage and the herbicide's penetration of weed foliage. See **Cropping Restrictions** for specified interval between application and planting to prevent crop injury.

For best performance, make application when annual weeds are less than 6" tall, when biennial weeds are in the rosette stage and to perennial weed regrowth in late summer or fall following a mowing or tillage treatment. Most effective control of upright perennial broadleaf weeds, such as Canada thistle and Jerusalem artichoke, occurs if application is made when the majority of weeds, such as field bindweed and hedge bindweed, are in or beyond the full bloom stage.

Avoid disturbing treated areas following application. Treatments may not kill weeds which develop from seed or underground plant parts, such as rhizomes or bulbets, after the effective period for **Disha DMA**. For seedling control, a follow-up program or other cultural practices could be instituted. For small grain in-crop uses of **Disha DMA**, refer to the **SMALL GRAINS** section.

Note: Observe all precautions. Read and follow mixing, application, and cleaning instructions.

Weed Type	Amount of Product per Treated Acre
Annual	½ - 1.0 pt.
Biennial	1.0 - 2.0 pts.
Perennial	
Suppression	1.0 - 2.0 pts.
Control - Noted with a superscript 1 in the following table.	2.0 pts.
Retreatments may be made as needed; however, do not exceed a total of 2 applications of Disha DMA per year.	

Disha DMA can be applied either post-harvest in the fall, spring, or summer during the fallow period or to crop stubble/set-a-side acres. **Disha DMA**, when applied at the specified rates, will control many annual broadleaf weeds, see the **Weeds Controlled** section under small grains. In addition, **Disha DMA** will control or suppress the following biennial and perennial broadleaf weeds:

Alfalfa ¹	Garlic, Wild ²	Spurge, Leafy
Artichoke, Jerusalem	Horsenettle, Carolina	Thistle, Bull
Bindweed, Field	Knapweed, Diffuse	Thistle, Canada ²
Bindweed, Hedge	Knapweed, Spotted	Thistle, Milk
Blueweed, Texas	Nightshade, Silver	Thistle, Musk
Bursage (Bur Ragweed, Povertyweed, Lakeweed) ¹	Redvine	Thistle, Plumeless
Dandelion, Common ¹	Smartweed, Swamp	Thistle, Scotch
Dock, Curly ¹	Sowthistle, Perennial ¹	Trumpet creeper (Buckvine)
Dogbone, Hemp		

¹These perennials may be controlled using **Disha DMA** at rates lower than those recommended for other listed perennial weeds.

²Refer to the **Special Tank Mix Treatments** section for specific control programs for these weeds.

Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Disha DMA may be applied alone or tank mixed with one or more of, but not limited to, the following herbicides for control of grasses or additional broadleaf weeds.

Broadcast Rate per Treated Acre For Annual Weed Control – Apply ¼ - 1.0 pt. (0.125 to 0.5 lb. a.e.) **Disha DMA** with:

Product	Active Ingredient	Formulation	Amount of Product per Acre
AAtrex® 4L ¹	atrazine	4.0 lbs./gal.	Refer to product label
AAtrex® Nine-O® ¹	atrazine	90% DF	Refer to product label
Ally® ²	metsulfuron-methyl	75% DF	Refer to product label
Amber® ²	triasulfuron	75% DF	Refer to product label
Fallow Master	glyphosate + dicamba	2.0 lbs./gal.	Refer to product label
Gramoxone® Extra	paraquat	2.5 lbs./gal.	Refer to product label
Kerb® ¹	pronamide	50-W	Refer to product label
Landmaster® BW	glyphosate + 2,4-D	2.4 lbs./gal.	Refer to product label
Roundup	glyphosate	3.0 lbs./gal.	Refer to product label
Sencor® DF ¹	metribuzin	75% DF	Refer to product label
Sencor® 4 ¹	metribuzin	4.0 lbs./gal.	Refer to product label
2,4-D	2,4-D	4.0 lbs./gal.	Refer to product label

¹Tank mixes of **Disha DMA** with these products may be subject to special restrictions. See the product label of the tank mix partner for intended use rates, restrictions. See the product label of the tank mix partner for intended use rates, restrictions, and other precautions.

²When tank mixing with sulfonylurea herbicides refer to the product label for rates and restrictions. Use a surfactant of at least 80% active ingredient at the rate of 1 - 2 qts./100 gals. of spray or not more than 0.25 - 0.5% by volume. Use the highest rate of surfactant when using the lower rate ranges of the tank mix and/or when treating more mature weeds or dense vegetative growth. Sulfonylurea-resistant weeds may not be controlled by tank mixes of **Disha DMA** and a sulfonylurea, refer to the **Disha DMA** tank mix section for alternative tank mixes.

Broadcast Rate per Treated Acre For Biennial and Perennial Weed Control – Apply 1.0 - 2.0 pts. (0.5 to 1 lb. a.e.) **Disha DMA** with:

Product	Active Ingredient	Formulation	Amount of Product per Acre
Curtail®	clopyralid + 2,4-D	2.38 lbs./gal.	Refer to product label
2,4-D	2,4-D	4.0 lbs./gal.	Refer to product label
Landmaster® BW	glyphosate + 2,4-D	2.4 lbs./gal.	Refer to product label
Roundup®	glyphosate	3.0 lbs./gal.	Refer to product label
Tordon® 22K	picloram	2.0 lbs./gal.	Refer to product label

Special Tank Mix Treatments

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

For suppression of perennial weeds, apply ½ - 1 pt. (0.25 to 0.5 lb. a.e.) **Disha DMA** with labeled use rate Roundup Herbicide or Roundup TR per treated acre.

For wild garlic control, apply 1 pt. (0.5 lb. a.e.) **Disha DMA** with labeled use rate of 2,4-D LV ester per treated acre. Apply when wild garlic is 4 - 8" tall.

For Canada thistle control, use **Disha DMA**, **Disha DMA** plus Curtail, or **Disha DMA** plus Roundup herbicide tank mix treatments.

Application may be made during fallow periods for control of volunteer barley, bulbous bluegrass, downy brome, jointed goat-grass, common rye, and volunteer wheat when they are actively growing. Use 1 pt. (0.5 lb. a.e.) **Disha DMA** with labeled use rate of Kerb 50-W. Fall seeded wheat may be planted 9 months or more after application. For best performance, make application between mid-October and mid-December, prior to soil freeze up.

During fallow periods, apply **Disha DMA** plus Landmaster BW or Fallow Master Herbicide to give improved control of Kochia, wild buckwheat, prickly lettuce, field bindweed, and Canada thistle. Use ¼ - ½ pt. (0.063 to 0.125 lb. a.e.) **Disha DMA** plus 22-54 fl. oz. of Landmaster BW or Fallow Master Herbicide for annual weed control or ¼ - ½ pt. (0.125 to 0.25 lb. a.e.) **Disha DMA** plus labeled use rate of Landmaster BW or Fallow Master Herbicide for perennial weed suppression.

Cropping Restrictions

The following restrictions are based on **Disha DMA** use rates up to 2 pts. (1 lb. a.e.) per treated acre:

- Corn, sorghum, and soybeans may be planted in the spring following applications made during the previous year. If less than 1" of rainfall occurs between application and the first killing frost, treated areas should be cultivated to allow herbicide to come in contact with moist soil. Cultivation may take place before or immediately after ground thaw.
- Soybean injury may occur if the interval between application and planting is less than specified. In areas with greater than 30" of rainfall, delay planting for 30 days per pt. of **Disha DMA** per treated acre. In areas with less than 30" of rainfall, delay planting for 45 days per pt. of **Disha DMA** per treated acre. Exclude days when ground is frozen.
- Wheat may be planted in the fall or spring following applications. also, spot applications may be made any time prior to crop emergence if crop injury can be tolerated in treated areas. Wheat injury may occur if the interval between application and planting is less than specified.
- East of the Mississippi River, the interval is 20 days per pt. of **Disha DMA** per treated acre or 1.25 days per 1 fl. oz. Moisture is essential for **Disha DMA** per treated acre or 3 days per fl. oz. Moisture is essential for **Disha DMA** degradation. Exclude days when ground is frozen.
- West of the Mississippi River, the interval is 45 days per pt. of **Disha DMA** per treated acre or 3 days per oz. Moisture is essential for **Disha DMA** degradation. Exclude days when ground is frozen.
- Following a normal harvest of barley, oats, or wheat, any rotational crop may be planted. If the interval before harvest is shortened, such as when cover crops will be plowed under, do not follow-up with the planting of a sensitive crop.

CONTROL OF PERENNIAL BROADLEAF WEEDS IN CROPLAND (SPOT APPLICATION* ONLY)

For Use Only in the States of Idaho, Montana, Nevada, Oregon, Utah, and Washington.

*Spot Application is defined as an area no greater than 1,000 ft. sq. per acre.

Disha DMA may be applied at any time following a crop harvest to stubble fallow or other cropland. Application should be made when weeds are actively growing and prior to a killing frost. Apply 1 qt. (1 lb. a.e.) of **Disha DMA** per treated acre. Application may be made up to 1 month prior to the planting of wheat.

Barley, oats, corn, sorghum (milo), annual or perennial grass crops may be planted into treated areas 1 year after application. Crops grown for seed (other than perennial grass seed) should not be planted into treated areas until 3 years after application. Do not plant broadleaf crops such as alfalfa, beans, peas, potatoes, or sugarbeets into treated areas until 2 years after application.

In most cases, treatments will not kill perennial weed seedlings which germinate from seed 1 or 2 years after treatment. Once the effect of the chemical has been lost, a follow-up program for seedling control or other cultural practices should be instituted.

Note: Observe all precautions. Read and follow mixing, application, and cleaning instructions.

Restrictions:

- Do not use unless injury to wheat or rotated barley will be acceptable.
- Do not treat sub-irrigated cropland or areas where the soil remains saturated with water throughout the year.
- Make only 1 application of **Disha DMA** per year.

Weeds Controlled

Disha DMA, when applied at specified rates, will control many broadleaf weeds including:

Bindweed, Field	Dock, Curly	Knapweed, Russian	Spurge, Leafy
Dock, Broadleaf (Bitterdock)	Knapweed, Black	Ragwort, Tansy	Thistle, Canada

Wiper Application Uses

Note: Observe all precautions. Read and follow mixing, application, and cleaning instructions.

Disha DMA may be applied through wiper application equipment to control or suppress actively growing broadleaf weeds, brush and vines. Use a solution contain 1 part **Disha DMA** to 1 part water. Do not contact desirable vegetation with herbicide solution. Wiper application should only be made to crops (including pastures) and non-cropland areas described in this label with the exception of grain sorghum (milo).

RIGHTS-OF-WAY, UTILITY AND INDUSTRIAL AREAS, AND FENCEROWS

Disha DMA is recommended for use on non-crop land areas such as rights-of-way (such as roadways, rest areas, utility, railroad, highway, pipeline, and rights-of-way that run through pasture and rangeland); utility facilities (such as substations, pipelines, tankfarms, pumping stations, parking and storage areas, fencerows, and non-irrigated ditchbanks); brush control for forest site preparation or maintenance.

Note: Observe all precautions on this label, including references to crops growing under stress, drift, sensitive crops, and tank cleanout.

- **Rights-of-Way - Disha DMA** can be used to control many broadleaf weeds on rights-of-way. This use includes applications to roadside, roadway and highways; to areas along utilities such as cable and powerlines; railroad track and embankment; highways, highway medians, bridge abutments, pipelines, and rights-of-way that run through pasture and rangeland. Use controlled application techniques that minimize the risk of off-target movement.
- **Utility and Industrial Areas - Disha DMA** can be used to control many broadleaf weeds and brush in non-crop areas on or surrounding substations, pipelines, tankfarms, pump stations, production facilities, and bareground situations. It may also be used on parking and storage areas.
- **Fencerows - Disha DMA** can be used to control many broadleaf weeds and brush in fencerows.

Tank Mixtures and Application

Read and observe **Management of Off-Site Movement** recommendations in this label.

Disha DMA can be applied using water, oil in water emulsions including invert systems, or sprayable fluid fertilizer as a carrier. A compatibility test (see **Compatibility Test** section) should be made prior to tank mixing.

To prepare oil in water emulsions, half-fill spray tank with water, then add the appropriate amount of emulsifier. With continuous agitation, slowly add the herbicide and then the herbicidal oil or a pre-mix of oil plus additional emulsifier to spray tank. Complete filling of spray tank with water. Maintain vigorous agitation during spray operation to prevent oil and water from forming separate layers.

Disha DMA may be applied broadcast using either ground or aerial application equipment. When using ground equipment, apply low or high volume sprays of between 3 - 600 gals. of diluted spray per treated acre. Volume of spray applied will depend on the height, density, and type of weeds or brush being treated and on the type of equipment being used. When using aerial equipment, apply 5 - 40 gals. of diluted spray per treated acre.

Disha DMA may be applied to individual clumps or small areas (spot treatment) of undesirable vegetation using handgun or similar types of application equipment. Apply diluted sprays to allow complete wetting (up to runoff) of foliage and stems.

Herbicide adjuvants or other spray additives (emulsifiers, spreader stickers, surfactants, wetting agents, drift control agents, or penetrants) may be used for wetting, penetration, or drift control. Spray additives must be agriculturally approved when used in pasture applications. If spray additives are used, read and follow all use recommendations and precautions on product label.

Weeds and Brush Controlled

Disha DMA, when applied at specified rates, will give control of many annual, biennial, and perennial broadleaf weeds, and many woody brush and vine species commonly found in non-crop land areas. Noted perennial weeds (superscript 1 in **WEED LIST** table) may be controlled with lower rates of either **Disha DMA** or **Disha DMA** plus tank mix combinations. Refer to the below **Rates and Timings** table.

Rates and Timings

Application rates and timings of **Disha DMA** are given below. Use the higher level of listed rate ranges when treating dense or tall vegetative growth.

Weed Stage & Type	Amount of Product per Acre	Gals. of Spray Mixture per Acre ²	Spray Concentration For Low Volume Application ⁴ (% Vol./Vol.)
Annual			
Small, Actively growing	½ - 1 pt.	25 - 50	3
Established weed growth	1 - 1 ½ pts.	50 - 75	3
Biennial¹ (Rosette diameter)			
Less than 3"	½ - 1 pt.	25 - 50	3 - 4
3" or more	1 - 2 pts.	50 - 100	3 - 4
Bolting	2 pts.	100 - 150	3 - 4
Perennial			
Suppression or top growth control	½ - 1 qt.	50 - 100	4
Noted Perennials (superscript 1 in WEED LIST table)	1 qt.	100 - 200	4
Other Perennials	1 qt.	200	5
Woody Brush and Vines³			
Top Growth Stems	½ - 1 qt.	50 - 200	5
and Roots	1 qt.	200	5

¹For best performance, make application when biennial weeds are in the rosette stage.

²Assuming typical application rate of 1 qt. of **Disha DMA** per 100 gals.

³Tank mixes may be required for optimal control. Refer to the **WEED LIST** table.

⁴Low volume rates must not exceed 4 pts. of **Disha DMA** maximum per acre per year (5% v/v = 10 gals. maximum solution per acre per year).

Retreatments may be made as needed; however, do not exceed 2 applications of **Disha DMA** per acre per year.

Tank Mix Options for Rights-Of-Way, Utility and Industrial Areas, and Fencerows

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Disha DMA may be tank mixed with other herbicides for additional weed control. Due to the differences that may occur between specific formulated products and specific use ingredients (e.g., water supplies), a compatibility test (see **Compatibility Test** section) is recommended prior to actual tank mixing. The following table lists example options, but does not limit tank mix options. Consult product labels for rate recommendations for tank mix partners.

Herbicide	Rates per Treated Acre
norflurazon (Predict)	Consult product labels for use rate and restrictions.
prodiamine (Endurance)	
glufosinate (Finale)	
glyphosate (Roundup, Accord)	
metsulfuron-methyl (Escort)	
pendimethalin (Pendulum)	
triclopyr (Redeem, Garlon)	
clopyralid (Transline)	
bromacil (Hyvar)	
chlorsulfuron (Telar)	
diquat (Reward)	
simazine (Princep)	
diuron (Karmex)	
fosamine ammonium (Krenite)	
hexazinone (Velpar)	
imazapyr (Arsenal)	
imazethapyr (Plateau)	
sulfometuron-methyl (Oust)	
sulfosate (Touchdown)	
tebuthiuron (Spike)	
2,4-D	

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a well-ventilated area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING [Less Than 5 Gallons]: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONTAINER HANDLING [Greater Than 5 Gallons]: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

CONTAINER HANDLING [For Bulk and Mini-Bulk Containers]: Refillable container. Refill this container with pesticide only. Do not use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by State and local authorities.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such

use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

All trademarks are the property of their respective owners.