

GROUP

6

HERBICIDE

# Bashazon 5L

**FOR POST-EMERGENCE USE IN ALFALFA GROWN FOR SEED PRODUCTION, BEANS, CLOVER  
GROWN FOR SEED, CORN, PEANUT, PEAS, PEPPERMINT, RICE, SORGHUM,  
SOYBEAN, AND SPEARMINT**

**ACTIVE INGREDIENT:**

Sodium salt of bentazon\* ..... 53.0%

**OTHER INGREDIENTS:** ..... 47.0%

**TOTAL:** ..... **100.0%**

\*Equivalent to 5.0 lbs. per gallon bentazon (3-(1-methylethyl)-1H,2,1,3-benzothiadiazin-4(3H)-one 2,2-dioxide) formulated as a soluble liquid.

**KEEP OUT OF REACH OF CHILDREN  
DANGER/PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

See label booklet for complete Precautionary Statements, Directions For Use, and Storage and Disposal.

**Manufactured For:**

**Sharda USA LLC** 

7217 Lancaster Pike, Suite A  
Hockessin, Delaware 19707

EPA Reg. No. 83529-66  
EPA Est. No. 70815-GA-002

**Net Contents: 2.5 Gallons**

<b>FIRST AID</b>	
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eyes open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after first 5 minutes; then continue rinsing eyes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give anything to an unconscious person.</li> </ul>
<b>HOTLINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at <b>1-800-222-1222</b> .	
<b>NOTE TO PHYSICIAN</b>	
Probable mucosal damage may contraindicate the use of gastric lavage.	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
DANGER**

Corrosive. Causes irreversible eye damage. DO NOT get in eyes or on clothing. Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear (goggles, face shield, safety glasses)

**USER SAFETY REQUIREMENTS**

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

**ENGINEERING CONTROLS**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

**IMPORTANT:** When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

## USER SAFETY RECOMMENDATIONS

### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. It is a violation of federal law to use any pesticide in a manner that results in the death of an endangered species or in adverse modification of their habitat.

## GROUNDWATER ADVISORY

Bentazon, the active ingredient in this product, is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This labeling must be in possession of the user at time of herbicide application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Observe all directions, precautions, and restrictions in this label and the labels of products used in combination with **Bashazon 5L** herbicide. Use of **Bashazon 5L** not consistent with this label can result in injury to crops, animals, or persons. Keep containers closed to avoid spills and contamination.

## PHYSICAL-CHEMICAL HAZARDS

Do not mix or allow coming in contact with oxidizing agent. Hazardous chemical reaction may occur.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

### Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, nitrile rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, or viton  $\geq 14$  mils
- Shoes plus socks
- Protective eyewear

## PRODUCT INFORMATION

**Bashazon 5L** provides selective post-emergence control of certain broadleaf weeds and sedges in alfalfa grown for seed, beans, clover grown for seed, corn, peanut, peas, peppermint, rice, sorghum, soybean, and spearmint. **Bashazon 5L** will not control grasses.

## USE RESTRICTIONS

- DO NOT apply more than 3.2 pints of **Bashazon 5L** per acre, per season in all crops.
- DO NOT apply more than 2.0 pounds of bentazon a.i. (from all sources) per acre, per season.

- DO NOT apply to stressed weeds that are subject to cold temperatures, lack moisture, or have herbicide injury, mechanical injury, as unsatisfactory control can result.
- DO NOT apply to crops growing in stressful conditions such as hail damage, flooding, drought, injury from other herbicides, or widely fluctuating temperatures, as crop injury can result.
- DO NOT apply to crops that show leaf phytotoxicity or plant stunting produced by previous herbicide applications as crop injury may be enhanced or prolonged.
- Rainfall or overhead irrigation within 4 hours after application reduces the effectiveness of **Bashazon 5L**.
- DO NOT apply through any type of irrigation system.
- DO NOT apply tank mixes in California.

### **Mode of Action**

**Bashazon 5L** is a Group 6 benzothiadiazinone herbicide and a photosynthesis PS II inhibitor. To ensure optimal performance, thoroughly cover target weeds with **Bashazon 5L**.

### **Crop Tolerance**

Crops listed in this label are tolerant to **Bashazon 5L**. Leaf speckling or bronzing can occur, but this condition is temporary; plants typically outgrow leaf speckling/bronzing within 10 days. **Bashazon 5L** does not interfere with new growth and crop vigor is maintained.

### **APPLICATION INSTRUCTIONS**

For optimal weed control, make post-emergent applications of **Bashazon 5L** to small, actively growing weeds early in the season. Make broadcast, band, or spot spray applications. Late applications allow weeds to exceed the maximum size and can prevent adequate control.

Apply **Bashazon 5L** at specified rates within the specified rate range to actively growing weeds prior to reaching the maximum sizes listed in **Tables 1, 3, and 4**.

If making sequential applications of **Bashazon 5L**, allow a minimum of 5 days between treatments.

### **Irrigation**

Ensure active weed growth in irrigated areas; as weeds stressed due to drought conditions are typically not adequately controlled.

### **Spray Coverage**

Ensure thorough spray coverage for optimum control of emerged weeds. Dense leaf canopies, shelter, and smaller weeds can prohibit thorough spray coverage. Thorough spray coverage is easiest to obtain early in the season when weeds are small.

### **Cultivation**

DO NOT cultivate within 5 days of application, or within 7 days of applying **Bashazon 5L**. Cultivation after 7 days may improve season-long weed control.

### **Application Methods and Equipment**

Avoiding spray drift at the application site is the responsibility of the applicator. The spray system and weather-related factors determine the potential for spray drift. The applicator is responsible for considering these factors when making application decisions to avoid spray drift onto non-target areas.

### **Ground Application**

Use a minimum spray volume of 10 gallons of water per acre to ensure adequate spray coverage. Use higher spray volume (up to 20 gallons of water per acre) to improve spray coverage when crop and weed foliage is dense. Apply using spray nozzles that deliver medium-to-coarse spray droplets as defined by ASABE standard S-572.1 and as shown in the nozzle manufacturer's spray catalogs. DO NOT use flood, whirl chamber, or controlled droplet applicator (CDA) nozzles or selective application equipment such as recirculating sprayers or wiper applicators. DO NOT use brass nozzles because of the corrosive effects of nitrogen additives.

### **Aerial Application**

Use a minimum spray volume of 5 gallons of water per acre. To obtain uniform coverage and to avoid off-target drift movement, applicators must observe the following requirements:

- **Boom Length** - The distance of the outermost nozzles on the boom must not exceed  $\frac{3}{4}$  the length of the fixed wingspan or 90% of rotor-blade diameter.
- **Nozzle Type** - Use low-drift nozzles such as straight stream (D-8 or larger). DO NOT use nozzles producing a mist droplet spray.
- **Nozzle Orientation** - Nozzles must always point backward parallel with the airstream and never be pointed downward more than 45 degrees.
- **Application Height** - Without compromising aircraft safety, apply at a height of 10 feet or less above the crop canopy or tallest plants.

## Aerial Application Directions

To obtain uniform coverage and to avoid drift hazards:

- **DO NOT** apply **Bashazon 5L** by aircraft when winds exceed 10 mph.
- Use coarse sprays (larger droplets) as they are less likely to drift.
- **DO NOT** apply **Bashazon 5L** by air if sensitive species (such as cotton, sugar beet, sunflower, or okra) are within 200 feet downwind.

**Table 1. Application Rates and Weeds Controlled in All Crops Except Rice\***

Weeds Controlled (includes ALS-resistant, glyphosate- resistant, and triazine-resistant biotypes)	Bashazon 5L Application Rates for Weed Growth Stages (Pints/A)					
	0.8		1.2		1.6	
	Weed Leaf Stage	Maximum Height (Inches)	Weed Leaf Stage	Maximum Height (Inches)	Weed Leaf Stage	Maximum Height (Inches)
Anoda, spurred	—	—	Up to 6	3	6 - 8	4
Balloonvine	—	—	2 - 4	2	4 - 6	3
Beggarticks	—	—	Up to 6	6	6 - 8	8
Bindweed, field <sup>8</sup>	See the <b>Special Directions for Problem Weeds</b> section.					
Bindweed, hedge <sup>8</sup>	See the <b>Special Directions for Problem Weeds</b> section.					
Buckwheat, wild	—	—	Up to 4	3	4 - 6	5
Canada thistle <sup>8</sup>	See the <b>Special Directions for Problem Weeds</b> section.					
Cocklebur <sup>1,8</sup>	2 - 4	4	2 - 6	6	6 - 10	10
Croton, tropic	—	—	Up to 2	2	2 - 4	4
Dayflower	—	—	Up to 6	4	6 - 10	8
Devils claw <sup>2</sup>	—	—	—	—	Up to 6	3
Eclipta	—	—	Up to 6	2	Up to 6	2
Galinsoga <sup>2</sup>	—	—	—	—	Cotyledon to 6	2
Groundsel, common	—	—	—	—	—	3
Jimsonweed	Up to 4	4	Up to 6	6	6 - 10	10
Ladysthumb	Up to 4	4	Up to 6	6	6 - 10	10
Lambsquarters, common <sup>2,3</sup>	Up to 4	1	Up to 6	1.5	Up to 6	2
Marshelder	—	—	Up to 4	2	Up to 8	4
Mayweed/Dogfennel	—	—	—	2	—	3

\*For **Bashazon 5L** use rates and weeds controlled in rice, refer to **Table 3. Application Rates and Weeds Controlled in Rice - Flooded Fields** and **Table 4.**

**Application Rates and Weeds Controlled in Rice - Drained Fields** in the **RICE** section.

<sup>1</sup>DO NOT treat earlier than leaf stage shown, and DO NOT count cotyledon leaves.

<sup>2</sup>Use crop oil concentrate (COC) or COC plus urea ammonium nitrate (UAN).

<sup>3</sup>For regrowth or new germination, a second application of **Bashazon 5L** may be necessary.

<sup>8</sup>See the **Special Directions for Problem Weeds** section.

(continued)

**Table 1. Application Rates and Weeds Controlled in All Crops Except Rice\* (cont.)**

Weeds Controlled (includes ALS-resistant, glyphosate- resistant, and triazine-resistant biotypes)	Bashazon 5L Application Rates for Weed Growth Stages (Pints/A)					
	0.8		1.2		1.6	
	Weed Leaf Stage	Maximum Height (Inches)	Weed Leaf Stage	Maximum Height (Inches)	Weed Leaf Stage	Maximum Height (Inches)
Morningglory <sup>5</sup>	—	—	4	4	6	6
Morningglory, cypressvine <sup>5</sup>	—	—	4	4	4	4
Morningglory, smallflower <sup>5</sup>	—	—	4	4	4	4
Mustard, wild	Up to 4	2	Up to 6	4	6 - 10	8
Nightshade, hairy <sup>7</sup>	—	—	—	—	2 - 6	4
Nutsedge, yellow <sup>8</sup>	See the <b>Special Directions for Problem Weeds</b> section.					
Poinsettia, wild <sup>2</sup>	—	—	Up to 6	4	4 - 8	6
Purslane, common	—	—	Up to 4	1	4 - 6	2
Radish, volunteer	—	—	2 - 6	4	6 - 10	10
Ragweed, common <sup>2</sup>	—	—	—	—	4 - 6	3
Ragweed, giant <sup>3</sup>	—	—	—	—	Up to 4	6
Redweed	—	—	4 - 6	6	6 - 10	8
Senna, coffee <sup>2</sup>	—	—	—	—	Up to 1 pinnate	2
Sesbania <sup>2</sup>	—	—	—	—	3 - 5	3
Shepherd's purse <sup>4</sup>	—	—	Up to 6	4	6 - 10	8
Sida, prickly (Teaweed)	—	—	Up to 6	3	6 - 8	4
Smartweed, Pennsylvania	Up to 4	4	Up to 6	6	6 - 10	10

\*For **Bashazon 5L** use rates and weeds controlled in rice, refer to **Table 3. Application Rates and Weeds Controlled in Rice - Flooded Fields** and **Table 4. Application Rates and Weeds Controlled in Rice - Drained Fields** in the **RICE** section.

<sup>2</sup>Use crop oil concentrate (COC) or COC plus urea ammonium nitrate (UAN).

<sup>3</sup>For regrowth or new germination, a second application of **Bashazon 5L** may be necessary.

<sup>4</sup>DO NOT treat rosette before seed stalk appears.

<sup>5</sup>Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, and Virginia - Make a second application 5 to 14 days later. For all other states, apply 1.6 to 2.4 pints of **Bashazon 5L** per acre to annual morningglory plants not larger than 4 true leaves. Control may be partial or inconsistent.

<sup>7</sup>**Bashazon 5L** does not control black nightshade or Eastern black nightshade.

<sup>8</sup>See the **Special Directions for Problem Weeds** section.

(continued)

**Table 1. Application Rates and Weeds Controlled in All Crops Except Rice\* (cont.)**

Weeds Controlled (includes ALS-resistant, glyphosate- resistant, and triazine-resistant biotypes)	Bashazon 5L Application Rates for Weed Growth Stages (Pints/A)					
	0.8		1.2		1.6	
	Weed Leaf Stage	Maximum Height (Inches)	Weed Leaf Stage	Maximum Height (Inches)	Weed Leaf Stage	Maximum Height (Inches)
Starbur, bristly	—	—	Up to 4	2	4 - 6	3
Sugar beet, volunteer	—	—	2 - 4	—	4 - 8	—
Sunflower, wild	Up to 2	3	Up to 4	5	4 - 6	8
Velvetleaf <sup>6,8</sup>	Up to 4	2	Up to 4	2	4 - 6	5
Venice mallow	Up to 4	2	Up to 6	2	6 - 10	4

\*For **Bashazon 5L** use rates and weeds controlled in rice, refer to **Table 3. Application Rates and Weeds Controlled in Rice - Flooded Fields** and **Table 4. Application Rates and Weeds Controlled in Rice - Drained Fields** in the **RICE** section.  
<sup>6</sup>Always use UAN or ammonium sulfate (AMS) as spray additive.  
<sup>8</sup>See the **Special Directions for Problem Weeds** section.

**Special Directions for Problem Weeds**

Weed Species	Bashazon 5L Application Rate & Timing	Special Instructions
<b>Field and Hedge Bindweed</b>	1.6 to 2.4 pints per acre when weeds smaller than 10 inches.	Suppression only in Kentucky, Illinois, Indiana, Michigan, and Ohio.
<b>Canada Thistle</b>	1.6 pints per acre when weeds smaller than 8 inches to bud stage.	Make a second application at the same rate 7-10 days later if regrowth occurs.
<b>Yellow Nutsedge</b>	1.2 to 1.6 pints per acre when weeds smaller than 8 inches.	Make a second application at the same rate 7-10 days later if regrowth occurs.
<b>Late Rescue of Cocklebur</b>	1.6 to 2.4 pints per acre when weeds smaller than 8 inches.	For optimal control, make an initial application of 1.2 pints per acre, then repeat application 10 to 14 days later.
<b>Late Rescue of Velvetleaf</b>	2.4 pints per acre plus spray additives (an oil concentrate and UAN, refer to <b>ADDITIVES</b> section for more details) when weeds are less than 12-inches tall.	For optimal control, make an initial application of 1.2 pints per acre plus spray additives (an oil concentrate and UAN or AMS, refer to <b>ADDITIVES</b> section for more details), then repeat application 5 to 7 days later.

## ADDITIVES

Apply **Bashazon 5L** with one or a combination of the additives listed below for optimal weed control. Additives must be added to the spray tank. The use of additives can cause some leaf burn, but new growth is normal and crop vigor doesn't change. Leaf burn commonly occurs in relatively humidity and high temperatures. When an adjuvant (or a specific adjuvant product, such as a drift control agent) is used with **Bashazon 5L**, use a Chemical Producers and Distributors Association (CPDA) certified adjuvant. See **Table 2. Additive Rates**. Applicators must use the additive(s) that will provide the best performance suited for their geography, target weed, and environmental conditions.

<b>Table 2. Additive Rates</b>	
<b>Additive/Adjuvant</b>	<b>Rate (Per Volume of Spray Solution)</b>
Crop oil concentrate (COC) <sup>1</sup>	1 gal./100 gals. (1% volume/volume [v/v])
Methylated seed oil concentrate (MSO) <sup>1,2</sup> (MSO adjuvant must contain at least 60% methylated seed oil. Poor performance may occur with adjuvants containing less than 60% methylated seed oil.)	1 gal./100 gals. (1% v/v)
Nonionic surfactant (NIS) (Use a NIS containing at least 80% active ingredient. Organosilicone surfactant may be used in place of NIS.)	1 - 2 qts./100 gals. (0.25% to 0.50% v/v)
<b>Additive/Nitrogen Fertilizer</b>	<b>Rate (Per Volume of Spray Solution)</b>
Ammonium sulfate (AMS) (use only spray-grade dry AMS)	8.5 - 17 lbs./100 gals. (1% to 2% weight/volume [w/v])
Urea ammonium nitrate (UAN) <sup>3</sup> (recommended liquid fertilizers include 28% N, 32% N, or 10-34-0)	1.25 - 2.5 gals./100 gals. (1.25% to 2.5% v/v)
<p><sup>1</sup>Petroleum-based or vegetable seed-based crop oil concentrate may be used. The oil concentrate must be nonphytotoxic, contain only EPA exempt ingredients, provide good mixing quality in a jar test, and be successful in local experience. The exact composition of suitable products will vary; however, petroleum-oil or vegetable-oil concentrates should contain emulsifiers to provide good mixing quality. Highly refined vegetable oils are more satisfactory than unrefined vegetable oils. To determine the suitability of oil concentrates with <b>Bashazon 5L</b>, conduct a jar test (see <b>Tank Mixing Information</b> section).</p> <p><sup>2</sup>MSO is recommended when weeds are under moisture or temperature stress.</p> <p><sup>3</sup>UAN may be added in place of other spray additives to improve control of cocklebur, devil's claw, Pennsylvania smartweed, velvetleaf, Venice mallow, wild mustard, and wild sunflower. <b>Bashazon 5L</b> plus a nitrogen solution will not provide adequate control of common ragweed and common lambsquarters. If these weeds or other weeds requiring oil concentrate are present in addition to velvetleaf, then oil concentrate should also be used.</p>	



## TANK MIXING INFORMATION

### DO NOT apply tank mixes in California.

Additives and/or other pesticides may be mixed in the spray tank with **Bashazon 5L** according to the specific tank mixing instructions in this label and respective product labels.

See Crop-specific Information for more details. Read and follow the applicable Restrictions and Limitations and Directions For Use on all products involved in tank mixing. The most restrictive labeling applies to tank mixes.

Physical incompatibility, reduced weed control, or crop injury may result from mixing **Bashazon 5L** with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers not specified on this label.

### Compatibility Test with Oil Concentrate

Before mixing components, always perform a compatibility jar test. Use only water from the intended source at the source temperature. For a spray volume of 20 gallons per acre, use 3.3 cups (800 mL) of water. For other spray volumes, adjust proportionately. Add 2 teaspoons each of herbicide and oil concentrate for each 1.6 pints per acre of label rate.

### Add components in the following sequence, gently mixing between additions:

1. **Bashazon 5L**
2. Tank mix product, if used
3. Oil concentrate

Cap jar, invert 10 cycles, let stand for 15 minutes.

**Evaluate.** An ideal tank mix combination will be uniform. The suitability of the oil concentrate is questionable if any of the following are observed:

- **Free oil at the surface** - Film or globules
- **Flocculation** - Fine particles which may be suspended in the liquid or found as a precipitated layer at the bottom of the jar.
- **Clabbering** - Thickening texture (coagulated) resembling yogurt or a curd-like texture as with cottage cheese.

### Mixing Order

Maintain constant agitation throughout mixing and application until spraying is complete.

1. Water - Fill tank ½ to full with clean water and start agitation.
2. Inductor - If an inductor is used, rinse it thoroughly after each component has been added.
3. Products in PVA bags - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
4. Water-soluble additives (including NIS, and nitrogen fertilizers such as AMS or UAN)
5. Water-dispersible products (such as dry flowables, wettable powders, suspension concentrates, or suspo-emulsions)
6. Water-soluble products (such as **Bashazon 5L**)
7. Emulsifiable concentrates (such as COC or MSO)
8. Remaining quantity of water.

### Cleaning Spray Equipment

Clean application equipment thoroughly by using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions, followed by triple rinsing the equipment before and after applying **Bashazon 5L**.

---

## CROP SPECIFIC DIRECTIONS

---

Follow all product information, mixing, application, weeds controlled, and additive instructions in the preceding sections of the label. Always read and follow all label directions when using this product alone, or when applying **Bashazon 5L** in tank mix combinations. The most restrictive labeling applies when using tank mixes.

## ALFALFA Grown for Seed

### For use ONLY in Idaho, Montana, Nevada, Oregon, Washington, and Wyoming.

Alfalfa seed is tolerant to post-emergence applications of **Bashazon 5L**. Leaf speckling, leaf bleaching or whitening, and temporary stunting can occur under certain conditions. Applications of **Bashazon 5L** made at or after flower bud formation can result in reduced seed yields.

Make broadcast post-emergence applications of **Bashazon 5L** in the spring. Apply 1.6 pints per acre **Bashazon 5L** to alfalfa with at least 2 trifoliolate leaves, but with no flower bud formation. For optimal control of troublesome weeds, make a second application of 1.6 pints per acre 5 to 14 days later.

Adding an oil concentrate (see **ADDITIVES** section for details) to **Bashazon 5L** on alfalfa grown for seed can cause slight leaf burn and/or temporary stunting, but new growth will be normal. Leaf burn is common in relatively high humidity and high temperatures. There are some oil concentrates have cause excessive leaf burn. Refer to your supplier of **Bashazon 5L** for information concerning successful local experience before purchasing any oil concentrate.

#### RESTRICTIONS - Alfalfa Grown for Seed:

- DO NOT apply **Bashazon 5L** if alfalfa shows evidence of leaf phytotoxicity and/or plant stunting that may result from prior herbicide applications. Such injury may be enhanced and/or prolonged if **Bashazon 5L** is applied.
- For use only in fields of alfalfa grown for seed production.
- DO NOT use on alfalfa produced for livestock feed.
- DO NOT use **Bashazon 5L** treated alfalfa seed for sprouting.

**IMPORTANT:** The use of **Bashazon 5L** on alfalfa grown for seed is a non-feed/nonfood use. If the **Bashazon 5L** applicator of this pesticide is not the producer, the applicator must provide a copy of this label to the producer of the crop. Producers of this crop who use **Bashazon 5L**, or cause **Bashazon 5L** use on a field they operate, must provide a copy of this product label to the seed conditioner.

**Bashazon 5L** does not have an established pesticide residue tolerance for alfalfa. Therefore, the **alfalfa seed crop, including forage or stubble, green chop, hay, pellets, meal, whole seed, cracked seed straw, or seed screenings**, cannot be used or distributed for human food or animal feed purposes.

Alfalfa seed treated with **Bashazon 5L** must bear a specific tag or conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading, with the following statement: "Not for human or animal consumption". All seed screenings from seed processing must be disposed of in such a manner that the screenings cannot be distributed or used for human food or animal feed purposes.

## BEANS, DRY and SUCCULENT

Adzuki, black turtle soup, cranberry, great northern, kidney, large lima, navy, pink, pinto, red, small lima, snap, and white beans

Not for use on adzuki beans in California.

Apply **Bashazon 5L** to beans after the first trifoliolate leaf has fully expanded. Even at the tolerant stages, yellowing, bronzing, speckling or burning of leaves is common under certain conditions, but is temporary and does not delay pod-set, maturity, or reduce yield. **Bashazon 5L** application with an oil concentrate (COC or MSO) can cause leaf burn, and may reduce yields, especially in snap bean.

**State-specific Instructions for Georgia and South Carolina:** DO NOT apply **Bashazon 5L** as a solo treatment to dry and succulent beans or severe crop injury may occur. Apply 4.8 to 12.8 fluid ounces of **Bashazon 5L** per acre to dry and succulent beans only when tank mixing with Raptor® or Pursuit® herbicides. Refer to the Raptor and Pursuit labels for additional use directions and restrictions. The most restrictive label directions apply.

#### RESTRICTIONS - Beans (Dry and Succulent):

- DO NOT apply to beans before the first trifoliolate leaf is fully expanded as severe crop injury may occur.
- DO NOT apply to garbanzo beans or lupines at any stage of growth.
- DO NOT apply within 30 days of harvest.

#### Tank Mixes

Tank mix **Bashazon 5L** in dry beans with one or more of the following herbicide products:

Clethodim

Outlook®

Poast®

Pursuit®

Raptor®

Reflex®

Other tank mix products may be used as long as the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

Tank mix **Bashazon 5L** in succulent beans with one or more of the following herbicide products:

Clethodim                      Poast®                      Pursuit®                      Reflex®                      Raptor® (snap bean only)\*

\*Only in states that Raptor is labeled for use on succulent lima bean and snap bean (see Raptor label).

Other tank mix products may be used as long as the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

### **CLOVER Grown for Seed For use ONLY in Oregon and Washington.**

Make post-emergence applications of **Bashazon 5L** to clover. Leaf burning is common under certain conditions, but plants generally outgrow this condition within 10 days. Apply **Bashazon 5L** with the appropriate additive and rate (see **ADDITIVES** section and **Table 2**).

Make a broadcast post-emergence application of 1.6 pints per acre of **Bashazon 5L** in the spring. To control troublesome weeds, make a second application of 1.6 pints per acre 5 to 14 days later.

#### **RESTRICTION - Clover Grown for Seed:**

- **Washington & Oregon:** DO NOT graze livestock or harvest forage or hay for livestock feed for at least 36 days after treatment.

#### **Tank Mixes**

**Bashazon 5L** may be applied in a tank mix with Raptor.

### **CORN**

#### **(field, sweet, pop, corn grown for seed or silage)**

Make post-emergence application of **Bashazon 5L**. Before making application to seed corn, verify the selectivity of **Bashazon 5L** on your inbred line with your local seed company (supplier) to avoid potential injury to sensitive inbreds.

#### **RESTRICTION - Corn:**

- DO NOT graze treated corn fields for at least 12 days after the last treatment with **Bashazon 5L**.

#### **Tank Mixes**

Tank mix **Bashazon 5L** with one or more of the following herbicide products:

Atrazine                      glyphosate (e.g., Roundup®)                      Outlook                      Status®

Other tank mix products may be used as long as the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

### **PEPPERMINT and SPEARMINT**

Application of **Bashazon 5L** to peppermint and spearmint can result in leaf burning when plants are actively growing and have extensive new, succulent tissue; however, plants generally outgrow this condition within 10 days.

#### **RESTRICTION - Peppermint and Spearmint:**

- DO NOT apply **Bashazon 5L** within 20 days of harvest.

#### **Tank Mixes**

Tank mix **Bashazon 5L** with one or more of the following herbicide products:

Buctril®                      Poast                      Sinbar®                      Stinger®

Other tank mix products may be used as long as the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

### **PEAS, DRY and SUCCULENT Including dry field, garden, English, and southern**

Make application of **Bashazon 5L** to peas after three pairs of leaves (or 4 nodes) are present. Yellowing, bronzing, speckling, or leaf burning is common under certain conditions, but injury does not delay pod-set, maturity, or reduce yield.

In-furrow application of insecticides or nematicides may predispose the peas to injury from **Bashazon 5L**.

**State-specific Instructions for Georgia and South Carolina:** Apply 4.8 to 12.8 fluid ounces per acre **Bashazon 5L** to dry and succulent peas only when tank mixed with Raptor® or Pursuit® herbicides. Refer to the Raptor and Pursuit labels for additional use directions or restrictions. DO NOT apply **Bashazon 5L** as a solo treatment to dry and succulent peas.

#### **RESTRICTIONS - Peas (Dry and Succulent):**

- DO NOT apply in western irrigated areas.
- DO NOT apply during prolonged periods (for 2 to 5 days) of cold weather (day temperature below 75°F and night temperature below 55°F).
- DO NOT apply to dry peas within 30 days or to succulent peas within 10 days of harvest in California.
- DO NOT apply to peas under stress from root rot.
- DO NOT apply to garbanzo beans or to lupines.

#### **Tank Mixes**

Tank mix **Bashazon 5L** in peas with one or more of the following herbicide products:

MCPA                      Pursuit                      Raptor                      Reflex®                      Thistrol®

Other tank mix products may be used as long as the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

Tank mixing of **Bashazon 5L** and Thistrol in the northeastern states of Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, Vermont, and in the mid-Atlantic states of Delaware, Maryland, Virginia, and in the Pacific northwestern states of Idaho, Oregon, and Washington:

Apply this tank mix after the 3-leaf stage (4-node stage) of peas, but not later than 3 nodes before pea flowering.

**NOTICE TO USER:** Because of variability among pea cultivars and in application techniques, neither the manufacturers nor the sellers have determined whether or not the tank mix of **Bashazon 5L** and Thistrol can be safely used on all pea crops under all conditions. Therefore, determine if the tank mix of **Bashazon 5L** and Thistrol can be used safely before large-scale use.

To improve control of pigweed and common lambsquarters, tank mix **Bashazon 5L** with MCPA.

#### **Tank Mix Restrictions and Limitations**

- DO NOT use oil concentrate, other oil-based additives, or any other spray additives or surfactants with these tank mixes.
- DO NOT apply the tank mix to peas when temperatures exceed 90°F.
- DO NOT apply the tank mix to peas after pea flower buds appear.
- Other crops, in particular beans, cotton, grapes, tomatoes, and ornamental plants, may be severely injured by off-target spray drift of Thistrol.

### **PEANUT**

Make application of **Bashazon 5L** from peanut cracking through pegging.

#### **RESTRICTIONS - Peanut:**

- DO NOT graze treated peanut fields for at least 50 days after the last **Bashazon 5L** treatment.
- Peanut hay and forage treated with **Bashazon 5L** may be fed to livestock.
- In-furrow treatments of insecticides and nematicides may predispose peanuts to injury from **Bashazon 5L**.

#### **Tank Mixes**

Tank mix **Bashazon 5L** with one or more of the following herbicide products:

2,4-DB amine    Cadre®    Cobra®    ET®    Gramoxone Inteon®    Outlook®    Poast®    Pursuit    Ultra Blazer®

Other tank mix products may be used as long as the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

### **Bashazon 5L and Gramoxone Inteon Tank Mix**

Apply this tank mixture at the ground crack stage to control an early flush of weeds. Make a second application up to 28 days after ground crack stage. Always add NIS at the specified rates. DO NOT use an oil concentrate or any other oil-based additive.

**Bashazon 5L** may be tank mixed with foliar fungicides such as Headline® SC fungicide and Priaxor® fungicide.

### **Peanut Tank Mix Restrictions and Limitations**

- DO NOT include UAN solution or AMS with the tank mix of **Bashazon 5L**, Poast® herbicide, and Ultra Blazer® herbicide.
- DO NOT add oil concentrate, UAN, or any other additives to the tank mix of **Bashazon 5L** and 2,4-DB.
- Use only amine formulations of 2,4-DB.
- Tank mixes not applicable in California.

## **RICE**

Make application of **Bashazon 5L** early post-emergence in rice, at the use rates specified below in and before weeds exceed the maximum size listed in **Tables 3 and 4**.

### **Alternate Flooding Culture**

**Arkansas, Louisiana, Mississippi, and Texas:** Weed growth stages generally correspond to rice that is tillering (stooling) and occur before the permanent flood. Apply **Bashazon 5L** when there is no water on the field and 24 hours or more before flooding. If **Bashazon 5L** cannot be applied until after flooding, see directions under **Continuous Flooding Culture**.

### **Continuous Flooding Culture**

In states using continuous flooding culture, or when treating after the permanent flooding, make application only when weeds are above the surface of the water. Weeds submerged at the time of application will not be adequately controlled. For early treatment, water may be partly or completely drained to expose more weed growth to spray applications of **Bashazon 5L**. DO NOT raise water level for at least 24 hours after application or unsatisfactory control may result. **DO NOT** use ground equipment to apply to flooded fields because splashing will wash **Bashazon 5L** off weed leaf surfaces and ineffective control may result.

Rice straw may be fed to livestock.

### **RESTRICTIONS - Rice:**

- DO NOT use **Bashazon 5L** on rice fields where commercial cultivation of catfish or crayfish is practiced.
- DO NOT use water containing **Bashazon 5L** residues from rice cultivation to irrigate crops used for food or feed unless **Bashazon 5L** is registered for use on these crops.
- DO NOT apply more than 3.2 pints of **Bashazon 5L** per acre per season whether one or two rice crops (including ratoon) are grown that season.

### **Tank Mixes**

Tank mix **Bashazon 5L** with one or more of the following herbicide products:

Beyond®	Command®	Facet® L	Grasp®	League™	Londax®	Ultra Blazer
Newpath®	Permit®	Permit Plus®	propanil	RiceBeaux®	Strada®	

Other tank mix products may be used, provided the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

### **Tank Mix Restrictions and Limitations**

- Apply the tank mix of **Bashazon 5L** and Londax within 7 days of establishing permanent flood.
- Apply the tank mix of **Bashazon 5L** and propanil only to drained fields.
- DO NOT use oil concentrate with the tank mix of **Bashazon 5L** and propanil. Add propanil to the tank mix of **Bashazon 5L** based on active ingredient (a.i.) of formulation used. Test propanil products for physical tank mix compatibility with **Bashazon 5L**.

### Special Directions for Problem Weeds in Rice

**Annual or Rice Flatsedge** (*Cyperus iria*) - For suppression, apply 1.2 to 1.6 pints of **Bashazon 5L** per acre when weeds are up to a maximum of 8-inches tall. If regrowth occurs, make a second application at the same rate 7 to 10 days later.

<b>Table 3. Application Rates and Weeds Controlled in Rice - Flooded Fields</b>				
<b>Weeds Controlled</b>	<b>Bashazon 5L Application Rates for Weed Growth Stages<sup>1</sup> (Pints/A)</b>			
	<b>1.2</b>		<b>1.6</b>	
	<b>Maximum Height Above Soil (Inches)</b>	<b>Height Range Above Water Level (Inches)</b>	<b>Maximum Height Above Soil (Inches)</b>	<b>Height Range Above Water Level (Inches)</b>
Cocklebur	10	3 - 6	15	6 - 10
Dayflower	6	3 - 5	10	5 - 8
Redstem	4	2 - 3	8	4 - 6
Smartweed	6	2 - 5	10	5 - 8
Water plantain, arrowhead	—	—	7	5 - 6
Water plantain, common	—	—	7	5 - 6
Yellow nutsedge	6	4 - 5	10	6 - 8

<sup>1</sup>If a second weed flush develops after the first application, re-treat according to this rate table.

<b>Table 4. Application Rates and Weeds Controlled in Rice - Drained Fields</b>				
<b>Weeds Controlled</b>	<b>Bashazon 5L Application Rates for Weed Growth Stages<sup>1</sup> (Pints/A)</b>			
	<b>1.2</b>		<b>1.6</b>	
	<b>Weed Leaf Stage</b>	<b>Maximum Height (Inches)</b>	<b>Weed Leaf Stage</b>	<b>Maximum Height (Inches)</b>
Cocklebur	2 - 10	10	10 - 15	15
Dayflower	2 - 10	6	10 - 15	10
Ducksalad	—	—	6 - 10	6
Eclipta	4 - 6	2	4 - 6	2
Gooseweed	4 - 6	4	6 - 10	8
Redstem	Up to 6	4	6 - 10	8
Redweed	4 - 6	6	6 - 10	8
Smartweed	2 - 10	6	10 - 15	10
Spikerush	2 - 6	6	6 - 8	8
Water plantain, arrowhead	—	—	Up to 4	7
Water plantain, common	—	—	Up to 4	7
Yellow nutsedge	4 - 6	6	6 - 8	10

<sup>1</sup>If a second weed flush develops after the first application, re-treat according to this rate table.

## RICE GROWN IN CALIFORNIA

Apply **Bashazon 5L** early post-emergence in rice at the use rates specified in **Table 5**.

Rice straw may be fed to livestock.

### Alternate Flood Culture

Weed growth stages generally correspond to rice that is tillering (stooling) and occur before the permanent flood. **Bashazon 5L** must be applied when there is no water on the field and 24 hours or more before flooding. If **Bashazon 5L** cannot be applied until after flooding, see directions under **Continuous Flood Culture**.

### Continuous Flood Culture

In continuous flood culture, or when treating after the permanent flooding, treatment must be made only when weeds are above the surface of the water. Weeds submerged at the time of application will not be adequately controlled. For early treatment, water may be partly or completely drained to expose more weed growth to spray applications of **Bashazon 5L**. DO NOT raise water level for at least 24 hours after application or unsatisfactory control may result. Application of **Bashazon 5L** with ground equipment to flooded fields can cause splashing that will wash **Bashazon 5L** off weed leaf surfaces and ineffective control may result.

### RESTRICTIONS - Rice Grown in California:

- DO NOT apply more than 3.2 pints of **Bashazon 5L** per acre per season.
- DO NOT release paddy water from field if **Bashazon 5L** has been applied during the growing season. Tail water may be recirculated within the field or other rice fields similarly treated with **Bashazon 5L**. Draining water from any **Bashazon 5L**-treated field during the growing season is not permitted.

### Tank Mixes

**Bashazon 5L** may be tank mixed in rice with, but not limited to, the following herbicide products:

Propanil

Other tank mix products may be used, provided the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

### Tank Mix Restrictions

- Apply the tank mix of **Bashazon 5L** and propanil only to drained fields.
- DO NOT use oil concentrate with the tank mix of **Bashazon 5L** and propanil. Add propanil to the tank mix of **Bashazon 5L** based on active ingredient (a.i.) of formulation used. Test propanil products for physical tank mix compatibility with **Bashazon 5L**.

Table 5. Application Rates and Weeds Controlled in Rice Grown in California				
Weeds Controlled	Bashazon 5L Application Rates for Weed Growth Stages <sup>1</sup> (Pints/A)			
	1.2		1.6	
	Application in Drained Fields Before Permanent Flooding			
	Weed Leaf Stage	Maximum Height (Inches)	Weed Leaf Stage	Maximum Height (Inches)
Ducksalad	—	—	6 - 10	6
Redstem	Up to 6	6	6 - 10	8

<sup>1</sup>If a second weed flush develops after the first application, re-treat according to this rate table.

(continued)

**Table 5. Application Rates and Weeds Controlled in Rice Grown in California (cont.)**

Weeds Controlled	Bashazon 5L Application Rates for Weed Growth Stages <sup>1</sup> (Pints/A)			
	1.2		1.6	
	Application to Flooded Fields			
	Maximum Height Above Soil (Inches)	Height Range Above Water Level (Inches)	Maximum Height Above Soil (Inches)	Height Range Above Water Level (Inches)
Redstem	4	2 - 3	8	4 - 6
Water plantain, common	—	—	7	5 - 6
Yellow nutsedge	6	4 - 5	10	6 - 8

<sup>1</sup>If a second weed flush develops after the first application, re-treat according to this rate table.

### SORGHUM, GRAIN and FORAGE

Make post-emergence application of **Bashazon 5L** to grain and forage sorghum. Before applying **Bashazon 5L** to sorghum, verify the selectivity of **Bashazon 5L** on your inbred or hybrid line with your local seed company (supplier) to help avoid potential injury to sensitive inbreds and hybrids.

#### RESTRICTIONS - Sorghum, Grain and Forage:

- DO NOT apply more than 1.6 pints of **Bashazon 5L** per acre per season.
- DO NOT apply to sorghum that is heading or blooming.
- DO NOT graze treated sorghum fields for at least 12 days after the last treatment with **Bashazon 5L**.

#### Tank Mixes

Tank mix **Bashazon 5L** in sorghum with one or more of the following herbicide products:

Atrazine                      Clarity®                      Facet® L                      Outlook®

Other tank mix products may be used as long as the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

### SOYBEAN

Make post-emergence applications of **Bashazon 5L** to soybean; soybean is tolerant to **Bashazon 5L** at all stages of growth; however, slight leaf speckling and leaf bronzing is common under certain conditions, but soybeans generally outgrow these conditions within 10 days.

#### RESTRICTION - Soybean:

- DO NOT graze or cut treated soybean fields for forage or hay for at least 30 days after the last treatment of **Bashazon 5L**.

#### Tank Mixes

Tank mix **Bashazon 5L** with one or more of the following herbicide products:

2,4-DB amine                      Cobra® herbicide                      Flexstar®                      glyphosate (e.g. Roundup® herbicide)                      Outlook  
 Poast®                      Pursuit®                      Raptor®                      Reflex®                      Resource®                      Ultra Blazer®

Other tank mix products may be used as long as the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

**Bashazon 5L** may be tank mixed with foliar fungicides and/or foliar insecticides (except malathion and Sevin® herbicide) if the application timings properly coincide.

#### Tank Mix Restrictions and Limitations

- **Bashazon 5L, Ultra Blazer, and Poast Tank Mixes:** Oil concentrate must be used with the tank mix of **Bashazon 5L**, Poast, and Ultra Blazer in place of a spray surfactant.



- **Bashazon 5L and 2,4-DB Amine Tank Mixes: Use** only amine formulations of 2,4-DB. Use no other adjuvant except UAN at 2 to 4 pints per acre with this tank mix. DO NOT apply more than 1 application of this tank mix per season. The use of this tank mix will cause soybean foliage injury (such as burning, bronzing, or crinkling) and may reduce yields. DO NOT use this tank mix on soybeans that show symptoms of disease such as *Phytophthora* root rot.
- The tank mixing of an insecticide with **Bashazon 5L** may increase the potential for crop injury.
- DO NOT apply tank mixes in California.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Do not allow this product to freeze.

**Pesticide Disposal:** Wastes resulting from this product may be disposed of on-site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

#### Container Handling:

**[Nonrefillable Container (five gallons or less):]** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

**[Nonrefillable Container (greater than five gallons):]** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

**[Refillable Container (greater than five gallons):]** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. Triple rinse as follows: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. When this container is empty, replace the cap and seal all openings that have been opened during use; return the container to the point of purchase or to a designated location. This container must only be refilled with a pesticide product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transport. DO NOT transport if this container is damaged or leaking. If the container is damaged, or leaking, or obsolete and not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling, if available, or dispose of container in compliance with state and local regulations.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.**

### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**BEFORE BUYING OR USING THIS PRODUCT,** read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of SHARDA USA LLC or the seller is authorized to vary in any way. Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product.

Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of SHARDA USA LLC and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, SHARDA USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS", AND SHARDA USA LLC MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT SHARDA USA LLC HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE MANUFACTURER.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF SHARDA USA LLC OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SHARDA USA LLC AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

All product names, brand names, and/or trademarks are property of their respective owners.

# BASHAZON 5L

**FOR POST-EMERGENCE USE IN ALFALFA GROWN FOR SEED PRODUCTION, BEANS, CLOVER GROWN FOR SEED, CORN, PEANUT, PEAS, PEPPERMINT, RICE, SORGHUM, SOYBEAN, AND SPEARMINT**

## ACTIVE INGREDIENT:

Sodium salt of bentazon\* ..... 53.0%

**OTHER INGREDIENTS:** ..... 47.0%

**TOTAL:** ..... 100.0%

\*Equivalent to 5.0 lbs. per gallon bentazon (3-(1-methylethyl)-1H,2,1,3-benzothiadiazin-4(3H)-one 2,2-dioxide) formulated as a soluble liquid.

## KEEP OUT OF REACH OF CHILDREN DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See label booklet for complete Precautionary Statements and Directions For Use.

### FIRST AID

**IF IN EYES:** Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes; then continue rinsing eyes. Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.

**HOTLINE NUMBER:** Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at **1-800-222-1222**.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Corrosive. Causes irreversible eye damage. DO NOT get in eyes or on clothing. Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **Pesticide Storage:** Do not allow this product to freeze. **Pesticide Disposal:** Wastes resulting from this product may be disposed of on-site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **Container Handling: [Nonrefillable Container (five gallons or less):]** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration. **[Nonrefillable Container (greater than five gallons):]** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration. **[Refillable Container (greater than five gallons):]** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. Triple rinse as follows: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. When this container is empty, replace the cap and seal all openings that have been opened during use; return the container to the point of purchase or to a designated location. This container must only be refilled with a pesticide product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transport. DO NOT transport if this container is damaged or leaking. If the container is damaged, or leaking, or obsolete and not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling, if available, or dispose of container in compliance with state and local regulations.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.**

**Manufactured For:** Sharda USA LLC, 7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707  
**EPA Reg. No. 83529-66      EPA Est. No. 70815-GA-002      Net Contents: 2.5 Gallons**