

GROUP 10 HERBICIDE

Opportunity

A non-selective herbicide for post-emergence broadcast use on canola, sweet corn*, corn, cotton, and soybean designated as LibertyLink®. Opportunity may be used for weed control in non-glufosinate tolerant cotton when applied with a hooded sprayer in-crop. Opportunity may also be applied as a broadcast burndown application before planting or before emergence of canola, sweet corn*, corn, soybean, and sugar beet* designated as LibertyLink® and any conventional canola, sweet corn*, corn, cotton, soybean, or sugar beet. Opportunity may be used for post-emergence weed control on olives, listed tree, vine, and berry crops. Opportunity may also be applied for potato vine desiccation.

*Not for use in California.

ACTIVE INGREDIENT:	WT. BY %
Glufosinate ammonium*24.5%**
OTHER INGREDIENTS:75.5%
TOTAL:	100.0%

*CAS Number 77182-82-2.

**Equivalent to 2.37 lbs. of active ingredient per U.S. gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

See label booklet for complete First Aid, Precautionary Statements, Directions For Use, and Storage and Disposal.

Manufactured For:

Sharda USA LLC 

7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

EPA Reg. No.: 83529-82

EPA Est. No.: 70815-GA-002, 39578-TX-001,
05905-IA-001, 228-IL-001

Net Contents: 2.5 Gallons

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to an unconscious person.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 .	
Note to Physician: If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible followed by charcoal and sodium sulfate administration.	

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton[®] ≥ 14 mils
- Shoes and socks
- Protective eyewear (goggles, face shield, or safety glasses)

All handlers must wear:

- Long-sleeved shirt and long pants
- Shoes and socks

Applicators using ground boom equipment with open cabs to treat cotton must wear:

- Long-sleeved shirt and long pants
- Shoes and socks
- Chemical-resistant gloves such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton[®] ≥ 14 mils

Mixer/loaders supporting ground boom applications to corn, canola, soybean, cotton, and olives must wear:

- Long-sleeved shirt and long pants
- Shoes and socks
- Chemical-resistant gloves such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton[®] ≥ 14 mils

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present. Do not apply to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters or rinsate.

This pesticide is toxic to vascular plants and must be used strictly in accordance with the drift and runoff precautions on this label in order to minimize off-site exposures.

Under some conditions, this product may have a potential to run off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no till, limited till and contour plowing. These methods also reduce pesticide run-off. Use of vegetation filter strips along rivers, creeks, streams, wetlands, etc., or on the downhill side of fields where run-off could occur to minimize water run-off is recommended.

PHYSICAL OR CHEMICAL HAZARDS

Do not use with or store near oxidizing agents since hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Not for Use in Nassau and Suffolk Counties in New York State.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

Exception: Sweet corn irrigation activities which has the restricted-entry interval (REI) of **4 days**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls worn over short-sleeved shirt and short pants
- Chemical-resistant gloves such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton[®] ≥ 14 mils
- Chemical-resistant footwear plus socks

IMPORTANT CROP SAFETY INFORMATION

READ BEFORE USING THIS PRODUCT

Opportunity may be applied as a burndown treatment before planting or before emergence of canola, sweet corn*, corn, soybean, and sugar beet* designated as LibertyLink[®] and any conventional canola, sweet corn*, corn, cotton, soybean, or sugar beet.

Post-emergence row crop applications of **Opportunity** may be made only to LibertyLink[®] crops.

The basis of selectivity of **Opportunity** in glufosinate-resistant crops is the presence of a gene tolerant to glufosinate. Crops not containing this gene will not be tolerant to **Opportunity** and severe crop injury and/or death may occur. Do not allow spray to contact foliage or green tissue of desirable vegetation other than LibertyLink[®] crops.

Opportunity may be applied to conventional or LibertyLink[®] cotton not tolerant to the active ingredient in **Opportunity** using a hooded sprayer.

Applications to trees, vines, and berries should avoid contact of **Opportunity** solution, spray drift, or mist with green bark, stems, or foliage, as injury may occur to trees, berries, and vines. Only trunks with callused, mature dark brown bark should be sprayed unless protected from spray contact by nonporous wraps, grow tubes or waxed containers. Contact of **Opportunity** with parts of trees, berries or vines other than mature brown bark can result in serious damage.

*Not for use in California.

RESISTANCE MANAGEMENT

Opportunity is classified as a Group 10 herbicide. Some pests are known to develop resistance to herbicides that have been used repeatedly. Pesticide resistance is the ability of a life form to develop a tolerance to a pesticide. While the development of weed resistance is well understood, it is not easily predicted. Therefore, herbicides should be used in conjunction with the resistance management strategies in the area. Consult the local or State agricultural advisors for details. If weed resistance should develop in the area, this product used alone may not continue to provide sufficient levels of pest control. If the reduced levels of control cannot be attributed to improper application techniques, improper use rates, improper application timing, unfavorable weather conditions or abnormally high pest pressure, a resistant strain may have developed.

To reduce the potential for pesticide resistance, use this product in a rotation program with other classes of chemistry and modes of action. Always apply this product at the specified rates and in accordance with the use directions. Do not use less than specified label rates alone or in tank mixtures. Do not use reduced rates of the tank mix partner. Users must scout before and after application and begin applications when pests are smaller rather than larger. If resistance is suspected, contact the local or State agricultural advisors.

INTEGRATED WEED PEST MANAGEMENT

Integrate **Opportunity** into an overall weed management strategy whenever the use of an herbicide is required. Practices known to reduce weed development (tillage, crop competition) and herbicide use (weed scouting, proper application timing, banding) should be followed wherever possible. Contact your local extension specialist, certified crop advisory and/or Sharda USA LLC representative for additional resistance management or IPM recommendation. Also, for more information of Weed Resistance Management, visit the Herbicide Resistance Action Committee (HRAC) on the web at: <http://www.hracglobal.com>

PRODUCT INFORMATION

Opportunity is a water soluble herbicide formulation for use as a foliar spray for the control of a broad spectrum of emerged annual and perennial grass and broadleaf weeds in canola, sweet corn*, corn, cotton, and soybean designated as LibertyLink®, and in olives, trees, vines, and berries. **Opportunity** may be applied for potato vine desiccation. **Opportunity** may also be applied as a broadcast burndown application before planting or before emergence of canola, sweet corn*, corn, soybean, and sugar beet* designated as LibertyLink® and any conventional canola, sweet corn*, corn, cotton, soybean, or sugar beet.

*Not for use in California.

Opportunity is only foliar active with little or no activity in soil. Weeds that emerge after application will not be controlled. Make application of **Opportunity** to actively growing weeds as described in the **WEED CONTROL FOR ROW CROPS** section to get maximum weed control. Uniform thorough spray coverage is necessary to achieve consistent weed control. Necrosis of leaves and young shoots occur within 2 to 4 days after application under good growing conditions.

- **Opportunity** is rainfast 4 hours after application to most weed species, therefore, rainfall within 4 hours may necessitate retreatment or may result in reduced weed control.
- Application should be made between dawn and 2 hours before sunset to avoid the possibility of reduced lambquarters and velvetleaf control.
- Consult your local Cooperative Extension Service or Sharda USA LLC representative for guidelines on the optimum application timing for **Opportunity** in your region.
- Weed control may be reduced if application is made when heavy dew, fog, and mist/rain are present, or when weeds are under stress due to environmental conditions such as drought, cool temperatures, or extended periods of cloudiness.

Restriction: To maximize weed control, do not cultivate from 5 days before an application to 7 days after an application.

SPRAY DRIFT MANAGEMENT

- When applying to crops via aerial application equipment, the spray boom must be mounted on the aircraft so as to minimize drift caused by wing tip or rotor blade vortices. The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter
- When applying to crops via aerial application equipment, applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.
- For aerial applications, do not release spray at a height greater than 10 feet above the crop canopy, unless a greater application height is required for pilot safety.
- For ground applications and aerial applications, select nozzle and pressure that deliver medium to coarse spray droplets as indicated in nozzle manufacturer's catalogues and in accordance with ASABE Standard 572.1.
- Spray at the appropriate boom height based on nozzle selection and nozzle spacing, but do not exceed a boom height of 24 inches above target pest or crop canopy. Set boom to lowest effective height over the target pest or crop canopy based on equipment manufacturer's directions. Automated boom height controllers are recommended with large booms to better maintain optimum nozzle to canopy height. Excessive boom height will increase the potential for spray drift.
- For non-crop vegetation management ground applications, apply with the nozzle height no more than 4 feet above the ground or target vegetation, unless necessitated by the application equipment. Examples would include roadside, railroad, utility rights of way, forestry and other industrial vegetation management applications where safety or natural barriers obstruct application.

SPRAY DRIFT ADVISORY

The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

POLLINATOR ADVISORY STATEMENT: This product contains an herbicide. Follow all label directions and precautions to minimize potential off-target exposure in order to prevent effects to non-target plants adjacent to the treated site which may serve as habitat or forage for pollinators.

Droplet Size Information

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS! (See **Wind, Temperature and Humidity**, and **Temperature Inversions**.)

Techniques for Controlling Droplet Size

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows usually produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Techniques for Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow Rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations. AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - Longer booms increase drift potential. Therefore, a shorter boom length is recommended.
- **Application Height** - Application more than 10 feet above the canopy increases the potential for spray drift.

Boom Height - Setting the boom at the lowest referenced height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

Drift Reduction Technology (DRT) - The EPA Drift Reduction Technology (DRT) Program was developed to encourage the manufacture, marketing, and use of spray technologies scientifically verified to significantly reduce pesticide drift. The use of DRTs should result in significantly less pesticide from spray applications drifting and being deposited in areas not targeted by those applications, compared to spray technologies that do not meet the minimum DRT standard. EPA-verified drift reduction technologies (DRTs) and their ratings will be added to the following webpage as they become available at: <https://www.epa.gov/reducing-pesticide-drift/epa-verified-and-rated-drift-reduction-technologies>

Wind - Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. AVOID APPLICATIONS DURING GUSTY OR WINDLESS CONDITIONS. **Note:** Local terrain can influence wind patterns. Every applicator needs to be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity - When making applications in hot and dry conditions, set-up equipment to produce larger droplets to reduce effects of evaporation.

Temperature Inversions - Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Shielded Sprayers - Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

APPLICATION AND MIXING PROCEDURES

Do not use flood jet nozzles, controlled droplet application equipment, or air assisted spray equipment. Uniform thorough spray coverage is important to achieve consistent weed control.

Ground Application

Refer to the Rate Tables for proper application rates. Do not apply when winds are gusty, or when conditions will favor movement of spray particles off the desired spray target. To avoid drift and ensure consistent weed control, apply **Opportunity** with the spray boom as low as possible while maintaining a uniform spray pattern. **Opportunity** should be applied broadcast in a minimum of 10 gallons of water per acre using a minimum spray pressure of 40 PSI and a maximum ground speed of 10 mph. The use of 80° or 110° flat fan nozzles is highly recommended for optimum spray coverage and canopy penetration. Application of the spray at a 45° angle forward will result in better spray coverage. **Under dense weed/crop canopies a broadcast rate of 15-20 gallons of water per acre should be used so that thorough spray coverage will be obtained.** DO NOT use raindrop nozzles. See the **SPRAY DRIFT MANAGEMENT** section of this label for additional information on proper application of **Opportunity**.

Aerial Application

Poor coverage will result in reduced weed control. For optimal weed control, apply **Opportunity** in a minimum of 10 gallons per acre. Apply **Opportunity** using nozzles and pressures that generate MEDIUM to COARSE spray droplets category as reported by the nozzle manufacturer and in accordance to ASABE S572 based upon the selected air speed. See the **SPRAY DRIFT MANAGEMENT** section of this label for additional information on proper application of **Opportunity**.

MIXING INSTRUCTIONS

Cleaning Instructions

Before using **Opportunity**, thoroughly clean bulk storage tank, refillable tank, nurse tanks, spray tank, lines, and filter, particularly if a herbicide with the potential to injure crops was previously used. Equipment should be thoroughly rinsed using a commercial tank cleaner.

After using **Opportunity**, triple rinse the spray equipment and clean with a commercial tank cleaner before using for crops not labeled LibertyLink®. Make sure any rinsate or foam is thoroughly removed from spray tank and boom. Rinsate may be disposed following the pesticide disposal directions on this label.

Tank Mixing Instructions

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Opportunity may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the crop to be treated. No label dosage rates may be exceeded. **Opportunity** cannot be mixed with any product containing a label prohibition against such mixing. Refer to the specific crop section for rates and restrictions. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Opportunity must be applied with properly calibrated and clean equipment. **Opportunity** is formulated to mix readily in water. Before adding **Opportunity** to the spray tank, ensure that the spray tank is thoroughly clean, particularly if a herbicide with the potential to injure crops was previously used (see **Cleaning Instructions**).

Mix **Opportunity** with water to make a finished spray solution as follows:

1. Fill the spray tank half full with water.
2. Start agitation.
3. If mixing with a flowable/wettable powder tank mix partner: Prepare a slurry of the proper amount of the product in a small amount of water. Add the slurry to the spray tank.
4. Add the appropriate amount of ammonium sulfate (AMS) to the spray tank.
5. If mixing with a liquid tank mix partner, add the liquid mix partner next.
6. Complete filling the spray tank with water.
7. Add the proper amount of **Opportunity** and continue agitation.
8. If foaming occurs, use a silicone based antifoam agent.

Ensure that all spray system lines including pipes, booms, etc., have the correct concentration of spray solution by flushing out the spray system lines before starting the crop application.

If tank mix partners specified on this label are added, maintain good agitation at all times until contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to re-suspend the mixture before spraying is resumed. Keep bypass line on or near bottom of tank to minimize foaming. Screen size in nozzles or line strainers must be 50-mesh or larger.

Compatibility Test for Tank Mixtures

If **Opportunity** is to be mixed with pesticide products not listed on this label, test the compatibility of the intended tank mixture before mixing the products in the spray tank. The following procedure assumes a spray volume of 25 gallons per acre. For other spray volumes, adjust the amount of the water used accordingly. Check compatibility as follows:

1. Place 1.0 pint of water from the source that will be used to prepare the spray solution in a clear 1 quart jar.
2. For each pound of dry tank mix partner to be applied per acre, add 1.5 teaspoons to the jar.
3. For each 16 fl. oz. of a liquid tank mix partner to be applied per acre, add 0.5 teaspoon to the jar.
4. For each 16 fl. oz. of **Opportunity** to be applied per acre, add 0.5 teaspoon to the jar.
5. After adding all the ingredients, place a lid on the jar and tighten. Invert 10 times to mix.
6. Let the mixture stand for 15 minutes, and evaluate the solution uniformity and stability. Look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. If the tank mix partners are not compatible, do not use the mixture in a spray tank.
7. After compatibility testing is complete, dispose of any pesticide wastes in accordance with the **Storage and Disposal** section of this label.

ROTATIONAL CROP RESTRICTIONS*

Rotational crop planting intervals following application of **Opportunity** are listed below. Failure to comply with these restrictions may result in illegal residues in rotated crops.

CROP	Minimum Rotation Interval (Days) After Last Opportunity Application
Canola, Sweet Corn, Corn, Cotton, Soybeans, and Sugar Beets	0 (May be planted at any time.)
Root and Tuber Vegetables, Leafy Vegetables, Brassica Leafy Vegetables, and Small Grains (barley, buckwheat, oats, rye, teosinte, triticale, and wheat)	70
All Other Crops	180

*See **DIRECTIONS FOR POTATO VINE DESICCATION** section for Rotational Crop Restrictions specifically after **Opportunity** applications to potatoes.

WEED CONTROL FOR ROW CROPS

Rates in ounces of formulated product per acre for the control of weeds at selected heights are shown in the weed control tables in weed populations with mixed species; make application at a rate needed for the species that requires the highest rate.

BROADLEAF WEED CONTROL

Weed Species	Maximum Weed Height or Diameter (Inches)		Weed Species	Maximum Weed Height or Diameter (Inches)	
	22 Fl. Oz./Acre	29 Fl. Oz./Acre ^{a,b}		22 Fl. Oz./Acre	29 Fl. Oz./Acre ^{a,b}
Amaranth, Palmer ²	NR	4	Morningglory, Smallflower ²	4	6
Anoda, Spurred	3	5	Morningglory, Tall ²	6	8
Beggarweed, Florida	4	5	Mustard, Wild	4	6
Black, Medic	5	7	Nightshade, Black	4	6
Blueweed, Texas	5	7	Nightshade, Eastern Black	6	8
Buckwheat, Wild	6	7	Nightshade, Hairy	6	8
Buffalobur	6	7	Pennycress (Stinkweed)	4	6
Burcucumber	6	10	Pigweed, Redroot ²	3	4
Catchweed Bedstraw (Cleavers)	2	4	Pigweed, Prostrate ²	3	4
Carpetweed	4	6	Pigweed, Spiny ²	3	4
Chickweed, Common	6	8	Pigweed, Smooth ²	3	4
Cocklebur, Common	6	14	Pigweed, Tumble ²	3	4
Copperleaf, Hophornbeam	4	6	Puncturevine	4	6
Cotton, Volunteer ¹	6	8	Purslane, Common	2	4
Croton, Tropic	3	5	Pusley, Florida	S	3
Croton, Woolly	2	4	Ragweed, Common	6	10
Eclipta	4	6	Ragweed, Giant	6	12
Devil's Claw	2	4	Senna, Coffee	4	6
Fleabane, Annual	6	8	Sesbania, Hemp	6	8
Galinsoga, Hairy	6	8	Shepherd's Purse	6	8
Galinsoga, Small Flower	6	7	Sicklepod (Java Bean)	4	6
Groundcherry, Cutleaf	4	5	Sida, Prickly	4	5
Geranium, Cutleaf	4	6	Smartweed, Pennsylvania	6	14
Hempnettle	4	6	Smellmelon	4	6
Horsenettle, Carolina ³	2	4	Sowthistle, Annual	6	8
Jimsonweed	6	10	Soybeans, Volunteer ¹	6	8
Knotweed	3	5	Spurge, Prostrate	2	4
Kochia ²	4	6	Spurge, Spotted	2	4
Ladysthumb	6	14	Starbur, Bristly	4	6
Lambsquarters, Common ²	4	6	Sunflower, Common	6	14
Mallow, Common	4	6	Sunflower, Prairie	3	5
Mallow, Venice	6	8	Sunflower, Volunteer	6	10
Marestail	S	6-12	Thistle, Russian ²	S	6-12
Marshelder, Annual	4	6	Velvetleaf ²	3	4
Morningglory, Entireleaf ²	6	8	Waterhemp, Common ²	NR	5
Morningglory, Ivyleaf ²	6	8	Waterhemp, Tall ²	NR	5
Morningglory, Pitted ²	6	8			
Morningglory, Sharpod ²	2	4			

^aIn cotton, **Opportunity** may be applied at 29 fl. oz./A, three times per year.

^b**Restriction:** Do not make application of more than 22 fl. oz./A of **Opportunity** post-emergence in a single application to canola and corn.

¹Volunteer LibertyLink crops from the previous season will not be controlled.

²For applications to corn, tank mixing with atrazine may enhance weed control of this species.

³May require sequential applications for control.

S = Indicates suppression.

NR = Not recommended.

GRASS WEED CONTROL

Weed Species	Maximum Weed Height or Diameter (Inches)		Weed Species	Maximum Weed Height or Diameter (Inches)	
	22 Fl. Oz./Acre	29 Fl. Oz./Acre ^{a,b}		22 Fl. Oz./Acre	29 Fl. Oz./Acre ^{a,b}
Barley, Volunteer ³	3	4	Millet, Wild Proso	6	7
Barnyardgrass	3	5	Millet, Proso Volunteer	6	7
Bluegrass, Annual	3	5	Oat, Wild ²	3	4
Corn, Volunteer ³	10	12	Panicum, Fall	3	5
Crabgrass, Large ²	3	5	Panicum, Texas	4	6
Crabgrass, Smooth ²	3	5	Rice, Red	4	6
Cupgrass, Woolly	6	12	Rice, Volunteer ¹	4	6
Foxtail, Bristly	6	8	Sandbur, Field ²	S	2
Foxtail, Giant	6	12	Shattercane	6	8
Foxtail, Green	6	12	Signalgrass, Broadleaf	3	5
Foxtail, Robust Purple	6	8	Sprangletop	4	6
Foxtail, Yellow ²	3	4	Sorghum, Volunteer	6	8
Goosegrass ³	2	3	Stinkgrass	4	6
Johnsongrass, Seedling	3	5	Wheat, Volunteer ²	4	5
Junglerice	3	5	Witchgrass	4	6

¹In cotton, **Opportunity** may be applied at 29 fl. oz./A, three times per year.

²**Restriction:** Do not make application of more than 22 fl. oz./A of **Opportunity** post-emergence in a single application to canola and corn.

³Volunteer LibertyLink crops from the previous season will not be controlled. A timely cultivation, 7 to 10 days after an application and/or retreatment for 10-21 days after the first application is recommended for controlling dense clumps of volunteer corn or rice.

²For best control of yellow foxtail, field sandbur, crabgrass, and wild oats, treat before initiation.

³A sequential application may be necessary for control.

S = Indicates suppression.

Biennial and Perennial Weeds**

For control of the biennial and perennial weeds listed below, tank mix partners or sequential applications of **Opportunity** are specified (22 fl. oz./A followed by 22 fl. oz./A).

Alfalfa	Clover, Alsike	Nutsedge, Purple*
Artichoke, Jerusalem	Clover, Red	Nutsedge, Yellow*
Bermudagrass	Dandelion	Orchardgrass
Bindweed, Field	Dock, Smooth	Poinsettia, Wild
Bindweed, Hedge	Dogbane, Hemp*	Pokeweed
Bluegrass, Kentucky	Goldenrod, Gray*	Quackgrass*
Blueweed, Texas	Johnsongrass, Rhizome	Sowthistle, Perennial
Bromegrass, Smooth	Milkweed, Common*	Thistle, Bull
Burdock	Milkweed, Honeyvine*	Thistle, Canada
Bursage, Woolyleaf	Muhly, Wirestem	Timothy*
Chickweed, Mouse Ear	Nightshade, Silverleaf	Wormwood, Biennial

*Suppression Only.

See the **DIRECTIONS FOR USE - COTTON section of this label for additional use rates.

APPLICATION DIRECTIONS FOR BURNDOWN USE

Opportunity may be applied as a burndown treatment before planting or before emergence of canola, sweet corn¹, corn, soybean, and sugar beet² designated as LibertyLink[®] and any conventional canola, sweet corn¹, corn, cotton, soybean, or sugar beet. Make application of a minimum of 29 fl. oz./A of **Opportunity** for burndown of existing weeds just before planting or before emergence of canola, corn, cotton, soybean, or sugar beets. For best results, make application to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of **Opportunity**. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures.

¹Not for use in California.

Crops	Application Directions
Canola, Sweet Corn, and Corn	If environmental conditions prevent timely applications, a single application may be made of up to 43 fl. oz./A of Opportunity . No additional applications of Opportunity may be made post-emergence to the crop during the growing season.
Cotton	If environmental conditions prevent timely applications, a single application may be made of up to 43 fl. oz./A of Opportunity . If more than 29 fl. oz./A are used in any single application, the yearly total may not exceed 72 fl. oz./A, including all application timings.
Soybean	If environmental conditions prevent timely applications, a single application may be made of up to 43 fl. oz./A of Opportunity . If 29-43 fl. oz./A are used in a single burndown application, one additional in season application may be made at up to 29 fl. oz./A. The yearly total may not exceed 87 fl. oz./A, including all application timings.
Sugar Beet*	If environmental conditions prevent timely applications, a single application may be made of up to 36 fl. oz./A of Opportunity . No additional applications of Opportunity may be made post-emergence to the crop during the growing season.
*Do not use on LibertyLink sugar beets in California.	

Crop (Conventional)	Burndown	In-Season Applications	Yearly Max
Cotton Use Pattern 1	29 fl. oz./A	2 applications at 29 fl. oz./A*	87 fl. oz./A
Cotton Use Pattern 2	30 - 43 fl. oz./A	1 application at 29 fl. oz./A*	72 fl. oz./A
Canola, Soybean, Sweet Corn, Corn	29 - 43 fl. oz./A	None	43 fl. oz./A
Sugar Beet	29 - 36 fl. oz./A	None	36 fl. oz./A
*Post-application in non-glyphosate cotton can only be applied with a hooded sprayer (see DIRECTIONS FOR USE - COTTON).			

Crop (LibertyLink® Varieties Only)	Burndown	In Season Applications	Yearly Max
Cotton Use Pattern 1	29 fl. oz./A	2 applications at 29 fl. oz./A	87 fl. oz./A
Cotton Use Pattern 2	30 - 43 fl. oz./A	1 application at 29 fl. oz./A	72 fl. oz./A
Canola	29 - 43 fl. oz./A	Up to 2 applications at 29 fl. oz./A	87 fl. oz./A
Sweet Corn, Corn, Soybean	29 - 43 fl. oz./A	Up to 2 applications at 29 - 43 fl. oz./A	87 fl. oz./A
Sugar Beet*	29 - 36 fl. oz./A	1 application at 29 fl. oz./A	60 fl. oz./A
*Not for use in California.			

DIRECTIONS FOR USE - SUGAR BEETS

(Not for use in California.)

THOROUGH SPRAY COVERAGE IS VERY IMPORTANT. **Opportunity** works best when weeds are actively growing. A cultivation may be made at least 5 days before an **Opportunity** application or 5 days after an **Opportunity** application.

Applications of **Opportunity** on sugar beets may be made from the cotyledon stage up to the 10-leaf stage of the sugar beet. **Opportunity** is a foliar active material with no soil residual activity. For best results, make application to emerged, young, actively growing weeds. Weeds that emerge after application will not be controlled. **Opportunity** will have an effect on weeds that are larger than the recommended leaf stage, however, speed of activity and control may be reduced. Weed control may be reduced if application is made when heavy dew, fog, and mist/rain are present, or when weeds are under stress due to drought, cool temperatures, or extended periods of cloudiness. **Opportunity** is rainfast 4 hours after application, therefore rainfall within 4 hours may necessitate retreatment.

For best weed control and sugar beet yield, **Opportunity** applications should begin when weeds are up to 1 inch in height or diameter. Repeat applications should be made when newly germinated weeds again reach 1 inch in height or diameter. Refer to the below **Rate Tables for Weed Control - Sugar Beets** for selection of the proper rate dependent upon the weed species present and size. A repeat application of **Opportunity** or a tank mix application with a residual herbicide selected from the tank mix partners listed on this label will be needed to control weeds that have not yet emerged at the time of application.

Restrictions - Sugar Beets

- Do not make application of more than 30 fl. oz./A (0.55 lb. a.i./A) of **Opportunity** in 1 application.
- Do not make application of more than 60 fl. oz./A (1.11 lbs. a.i./A) of **Opportunity** on the sugar beet crop per year.
- Do not apply more than 2 applications per year.
- Do not make application of **Opportunity** within 60 days of harvesting sugar beets.
- Do not plant rotation crops in a field treated with **Opportunity** within 120 days after the last application of this product with the exception of wheat, barley, buckwheat, millet, oats, rye, sorghum, and triticale, which may be planted 70 days after the last application of this product. LibertyLink corn, soybeans, canola, and sugar beets may be planted at any time.
- Do not graze the treated crop or cut for hay.
- Do not add surfactants. Antifoams or drift control agents may be added if needed.
- Do not make application of **Opportunity** if sugar beets show injury from prior herbicide or environmental stress (drought, excessive rainfall, etc.).
- Do not make application of **Opportunity** through any type of irrigation system.

Rate Tables for Weed Control - Sugar Beets

The rate of **Opportunity** in fluid ounces (pints) of formulated product per acre to be used for the control of weeds at selected heights is shown in the following tables. In weed populations with mixed species, make application at the rate needed for all species present.

Grass Weeds Controlled with Opportunity

Weed Species	Growth Stage of Weed* (Maximum Height)		Comments on Weed Growth Stage/Application Timing/Number of Applications
	15 Fl. Oz./Acre (0.9 Pt./A)	20 Fl. Oz./Acre (1.25 Pts./A)	
Barley, Volunteer	1 - 2 leaf (2")	3 leaf (3")	Multiple applications may be required
Barnyardgrass	1 - 3 leaf (2")	4 - 5 leaf (3")	Maximum of 1 tiller
Corn, Volunteer	1 - 2 leaf (3")	3 - 4 leaf (6")	-
Crabgrass, Large	1 - 3 leaf (2")	4 - 5 leaf (3")	Maximum of 1 tiller
Crabgrass, Smooth	1 - 3 leaf (2")	4 - 5 leaf (3")	Maximum of 1 tiller
Cupgrass, Woolly	1 - 5 leaf (4")	(8")	-
Foxtail, Giant	1 - 4 leaf (3")	5 - 6 leaf (4")	Maximum of 2 tillers
Foxtail, Green	1 - 4 leaf (3")	5 - 6 leaf (4")	Maximum of 2 tillers
Foxtail, Yellow	1 - 3 leaf (1")	4 leaf (2")	Make application before tillering
Millet, Volunteer Proso	1 - 3 leaf (2")	4 - 5 leaf (3")	Maximum of 1 tiller
Millet, Wild Proso	1 - 3 leaf (2")	4 - 5 leaf (3")	Maximum of 1 tiller
Oat, Wild	1 - 2 leaf (2")	3 leaf (3")	Maximum of 1 tiller
Panicum, Fall	1 - 3 leaf (2")	4 - 5 leaf (3")	Maximum of 1 tiller
Panicum, Texas	1 - 3 leaf (2")	4 - 5 leaf (3")	Maximum of 1 tiller
Sandbur, Field	-	1 - 4 leaf (2")	Make application before tillering
Wheat, Volunteer	1 - 2 leaf (2")	3 leaf (3")	Maximum of 1 tiller

*Make application up to 30 fl. oz./A (1.88 pts./A) if weeds exceed the growth stage shown in the table.

For improved control of heavy populations or larger than recommended volunteer wheat, volunteer barley, yellow foxtail, and wild oats, **Opportunity** can be tank mixed with Assure® II Herbicide, Poast® Herbicide, Prism® Herbicide, or Select® 2EC Herbicide.

Perennial Weeds Controlled by Opportunity

Weed Species	Growth Stage of Weed* (Maximum Height/Diameter)		Comments on Number of Applications
	15 Fl. Oz./Acre (0.9 Pt./A)	20 Fl. Oz./Acre (1.25 Pts./A)	
Quackgrass	-	1 - 3 leaf (3")	Multiple applications required
Sowthistle, Perennial	-	1 - 4 leaf (3")	Multiple applications required
Thistle, Canada	-	1 - 4 leaf (3")	Multiple applications required

*Make application up to 30 fl. oz./A (1.88 pts./A) if weeds exceed the growth stage shown in the table.

Broadleaf Weeds Controlled by Opportunity

Weed Species	Growth Stage of Weed* (Maximum Diameter)	
	15 Fl. Oz./Acre (0.9 Pt./A)	20 Fl. Oz./Acre (1.25 Pts./A)
Buckwheat, Wild	1 - 4 leaf (2")	5 - 6 leaf (3")
Buffalobur	1 - 4 leaf (2")	5 - 6 leaf (3")
Carpetweed	-	1 - 4 leaf (2")
Chickweed, Common	1 - 4 leaf (2")	5 - 6 leaf (3")
Cocklebur, Common	1 - 6 leaf (3")	7 - 8 leaf (5")
Kochia	(1")	(2")
Ladysthumb	1 - 2 leaf (1")	3 - 4 leaf (3")
Lambsquarters, Common	1 - 2 leaf (1")	4 - 5 leaf (3")
Mallow, Venice	1 - 4 leaf (2")	5 - 6 leaf (3")
Marshelder	1 - 2 leaf (1")	3 - 4 leaf (2")
Mustard, Wild	1 - 4 leaf (2")	5 - 6 leaf (3")
Nightshade, Eastern Black	1 - 4 leaf (2")	5 - 6 leaf (3")
Pigweed, Prostrate	(1")	(3")
Pigweed, Redroot	1 - 2 leaf (1")	3 - 4 leaf (3")
Pigweed, Smooth	1 - 2 leaf (1")	3 - 4 leaf (3")
Pigweed, Spiny	1 - 2 leaf (1")	3 - 4 leaf (3")
Purslane, Common	(1")	(2")
Ragweed, Common	1 - 6 leaf (3")	7 - 8 leaf (5")
Ragweed, Giant	1 - 4 leaf (2")	5 - 6 leaf (3")
Shepherd's Purse	1 - 4 leaf (2")	5 - 6 leaf (3")
Smartweed, Pennsylvania	1 - 2 leaf (1")	3 - 4 leaf (3")
Sowthistle, Annual	1 - 4 leaf (2")	5 - 6 leaf (3")
Sunflower, Common	1 - 6 leaf (3")	7 - 8 leaf (5")
Thistle, Russian	(1")	(2")
Velvetleaf	1 - 2 leaf (1")	3 - 4 leaf (3")

*Make application up to 30 fl. oz./A (1.88 pts./A) if weeds exceed growth stage shown in the table.

DIRECTIONS FOR USE - CANOLA

Make application of **Opportunity** only to canola labeled as LibertyLink. Uniform, thorough spray coverage is necessary to achieve consistent weed control. For best results, make application to emerged, young actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of **Opportunity**. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. For optimal yield, early season weed removal is important.

Applications of **Opportunity** on canola may be made from the cotyledon stage up to the early bolting stage of the canola. Slight discoloration of the canola may be visible after application. This effect is temporary and will not influence crop growth maturity or yield.

Make application of **Opportunity** at 22 - 29 fl. oz./A (0.41 - 0.54 lb. a.i./A) per application. A second application of **Opportunity** may be needed to control weeds that have not yet emerged at the time of application.

Restrictions - Canola

- Do not use on canola in the states of Alabama, Delaware, Georgia, Kentucky, Maryland, New Jersey, North Carolina, South Carolina, Tennessee, Virginia, and West Virginia.
- Do not apply more than 2 applications of **Opportunity** per year. Sequential applications need to be at least 7 days apart.
- Do not make application of **Opportunity** within 65 days of harvesting canola.
- Do not make application of more than 87 fl. oz. (1.61 lbs. a.i./A) of **Opportunity** per year.
- If **Opportunity** was used in a burndown application, no post-emergence applications may be applied to the crop.
- Do not graze the treated crop or cut for hay.
- Do not make application of **Opportunity** if canola shows injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
- Do not make application of **Opportunity** through any type of irrigation system.

Refer to the **ROTATIONAL CROP RESTRICTIONS** section for the appropriate rotational crop information.

Spray Additives

Opportunity must be applied with ammonium sulfate (AMS). Use only fine feed grade or spray grade AMS at 3 lbs. per acre. Anti-foams or drift control agents may be added if needed. Use of additional surfactants or crop oils may increase risk of crop response.

Tank Mixtures - Canola

Opportunity at 22 fl. oz./A plus AMS may be used in tank mix combination with certain herbicides for improved control of larger than labeled grasses. The AMS rate may be reduced to 1.5 lbs./A when **Opportunity** is tank mixed with a reduced rate of one of the grass herbicides specified below.

Opportunity may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the canola to be treated. When tank mixing, do not exceed specified application rates. **Opportunity** cannot be mixed with any product containing a label prohibition against such mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Tank Mix Partners for Opportunity on Invigor LibertyLink Canola

Tank Mix Partner	Rate (Fl. Oz./Acre)
Assure® II	Refer to product label
Poast®	Refer to product label
Select® ZEC	Refer to product label
Select Max™	Refer to product label

APPLICATION RATE AND TIMING TO LIBERTYLINK CANOLA FOR SEED PROPAGATION (Not for use in California.)

Up to 3 applications of **Opportunity** at up to 29 fl. oz./A per application may be made to LibertyLink canola for transgenic seed propagation. Applications may be made from the cotyledon stage up to the early bolting stage (e.g., BBCH 18-30, between just prior to stem elongation/bolting, eight or more leaves and beginning of stem elongation, no internodes).

Restrictions - LibertyLink Canola For Seed Propagation

- Do not make application of more than 3 applications of **Opportunity** at up to 29 fl. oz./A (0.54 lb. a.i./A) per application per year.
- Do not make application of more than 87 fl. oz./A (1.61 lbs. a.i./A) of **Opportunity** per year.
- Do not make application of **Opportunity** beyond the early bolting stage or within 65 days of harvesting canola seed.
- Do not use treated canola seed for food, feed, or oil purposes.
- Do not make application of **Opportunity** if canola shows injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
- Do not make application of **Opportunity** through any type of irrigation system.

DIRECTIONS FOR USE - SWEET CORN (Not for use in California.)

Make application of **Opportunity** only to sweet corn labeled as LibertyLink. Applications for **Opportunity** on sweet corn may be made from emergence until sweet corn is 24" tall or in the V-7 stage of growth (i.e., 7 developed collars), whichever comes first. Make application at a rate of 22 fl. oz./A. **Opportunity** must be applied with ammonium sulfate (AMS) for use on sweet corn. Two applications of **Opportunity** can be made to sweet corn in a year. Refer to the "ROTATIONAL CROP RESTRICTIONS" section for the appropriate rotational crop plant back intervals. See "DIRECTIONS FOR USE - FIELD CORN AND SILAGE CORN" for Application Methods, Mixing Instructions, and Weed Control Tables.

Restrictions - Sweet Corn

- Do not make application of **Opportunity** within 50 days of harvesting sweet corn ears and within 55 days of harvesting stover.
- Do not make application of more than 44 fl. oz./A (0.81 lb. a.i./A) of **Opportunity** on sweet corn per year.
- Do not make application of more than 2 applications of **Opportunity** to the sweet corn crop. Sequential applications need to be at least 7 days apart.
- If **Opportunity** was used in a burndown application, no post-emergence applications may be made to the crop.
- Do not use nitrogen solutions as spray carriers. A silicone-based antifoam agent may be added if needed.
- Do not make application of **Opportunity** if corn shows injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
- Do not make application of **Opportunity** through any type of irrigation system.

Tank Mixtures - Sweet Corn

Opportunity may be tank mixed with Laudis® Herbicide, Callisto®, Atrazine, or Permit®. When using **Opportunity** in tank mix combinations, carefully follow the "Directions for Use" labeling of the selected partner. When tank mixing, do not exceed specified application rates. **Opportunity** cannot be mixed with any product containing a label prohibition against such mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

DIRECTIONS FOR USE - FIELD CORN AND SILAGE CORN

Make application of **Opportunity** only to corn labeled LibertyLink. Uniform thorough spray coverage is necessary to achieve consistent weed control. For best results, make application to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of **Opportunity**. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. For optimal yield, early season weed removal is important.

Applications of **Opportunity** on corn may be made with over the top broadcast or drop nozzles from emergence until corn is 24" tall or in the V-7 stage of growth (i.e., 7 developed collars), whichever comes first. For corn 24" to 36" tall only, apply **Opportunity** using ground application and drop nozzles and avoid spraying into the whorl or leaf axils of the corn stalks. Applications of **Opportunity** following the use of soil applied insecticides will not injure corn.

Make application of **Opportunity** at 29 - 43 fl. oz./A (0.54 - 0.80 lb. a.i./A per application. A second application of **Opportunity** or a tank mix application with a residual herbicide will be needed to control weeds that have not yet emerged at the time of application.

Restrictions - Field Corn and Silage Corn

- Do not make application of **Opportunity** within 60 days of harvesting corn forage and within 70 days of harvesting corn grain and corn fodder.
- Do not make application of more than 2 applications of **Opportunity** to the corn crop per year. Sequential applications need to be at least 7 days apart.
- Do not make application of more than 87 fl. oz./A (1.61 lbs. a.i./A) of **Opportunity** on corn per year.
- If **Opportunity** was used in a burndown application, no post-emergence applications may be applied to the crop.
- Do not use nitrogen solutions as spray carriers. A silicone-based antifoam agent may be added if needed.
- Do not make application of **Opportunity** if corn shows injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
- Do not make application of **Opportunity** through any type of irrigation system.

Refer to the **ROTATIONAL CROP RESTRICTIONS** section for the appropriate rotational crop information.

Spray Additives

For corn and sweet corn, **Opportunity** must be applied with ammonium sulfate (AMS). It is recommended to use only fine feed grade or spray grade AMS at 3 lbs. per acre (17 lbs./100 gals.). When temperatures exceed 85°F, the rate of AMS can be reduced to 1.5 lbs. per acre (8.5 lbs./100 gals.) to reduce potential leaf burn. Use of additional surfactants or crop oils may increase risk of crop response.

Tank Mixtures - Corn

Certain herbicide tank mixes may aid in the performance of **Opportunity**. No additional surfactant is needed with any tank mix partner. **Opportunity** may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the corn to be treated. When tank mixing, do not exceed specified application rates. **Opportunity** cannot be mixed with any product containing a label prohibition against such mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Tank Mix Partners for Opportunity on LibertyLink Corn

2,4-D	Halex GT	Pendimethalin ¹
acetochlor	Hornet® WDG	Permit®
Aim™ ²	Impact®	Python® WDG
Atrazine	Laudis®	s metolachlor ²
Calisto™	Lexar® ²	Spirit®
Carmix® ²	Lumax® ²	Status®
Capreno®	Metolachlor ²	Yukon®
Distinct™	nicosulfuron	Zemax
Guardsman Max®	NorthStar™	

¹Tank mixing with pendimethalin may result in reduced control of barnyardgrass, fall panicum, field sandbur, yellow foxtail, and volunteer corn.

²It is directed that these products are tank mixed at half the use rate with **Opportunity** to reduce risk of crop response.

Corn Insecticide Tank Mix Partners for Opportunity

To provide weed and insect control in corn, **Opportunity** may be mixed with the following insecticides:

Ambush® Insecticide	Tombstone™ Helios®	Pounce® 3.2EC Insecticide
Asana® XL Insecticide	Lorsban® 4E Insecticide	Warrior™ Insecticide
Baythroid® XL Insecticide	Tombstone™	

DIRECTIONS FOR USE - COTTON

Uniform thorough spray coverage is necessary to achieve consistent weed control. **Opportunity** may be applied as a broadcast, over-the-top, post-emergence spray or as a directed spray only to LibertyLink cotton. **Opportunity** may be applied post-emergence to non-glufosinate tolerant cotton, varieties or cultivars by using equipment designed to minimize contact of the spray with the cotton foliage. Refer to the below **Non-Glufosinate Tolerant Cotton** section for selection of shielding equipment. Severe injury or death may result if the **Opportunity** contacts the foliage or stems of cotton NOT labeled as LibertyLink.

For best results, make application to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of **Opportunity**. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. For optimum yield, early season weed removal is important.

Make application of **Opportunity** to cotton from emergence up to the early bloom stage at 22 to 29 fl. oz./A. Should environmental conditions prevent a timely herbicide application, a single application of up to 43 fl. oz./A of **Opportunity** may be made to cotton. If more than 29 fl. oz./A are used in any single application, the yearly total may not exceed 72 fl. oz./A, including all application timings. See Restrictions to the **DIRECTIONS FOR USE - COTTON** below for additional information.

Refer to the **WEED CONTROL TABLE FOR ROW CROPS** section of this label for selection of the proper rate dependent upon weed species present and size. In weed populations with mixed species, select the highest rate required to control all the species. Volunteer LibertyLink crop plants (corn, cotton, soybeans, and sugar beets) from the previous season will not be controlled by applications of **Opportunity**. A repeat application of **Opportunity** or tank mixes with a residual herbicide will be needed to control weeds that have not emerged at the time of application. See the **Tank Mixtures - Cotton** section to select suitable tank mix partners.

Use Pattern	1 st Application	2 nd Application	3 rd Application	Yearly Maximum
Option 1	22 - 29 fl. oz./A	22 - 29 fl. oz./A	22 - 29 fl. oz./A	87 fl. oz./A
Option 2	30 - 43 fl. oz./A	22 - 29 fl. oz./A	None	72 fl. oz./A

Restrictions - Cotton

- Do not make application of **Opportunity** to cotton in Florida - South of Tampa (Florida Route 60), or in Hawaii (except for test plots or breeding nurseries).
- Do not make application of **Opportunity** within 70 days before cotton harvest.
- Up to 3 applications of **Opportunity** may be made to cotton per year at a maximum application rate of 29 fl. oz./A (0.54 lb. a.i./A). Sequential applications need to be at least 10 days apart.
- Do not make application of more than 87 fl. oz. (1.61 lbs. a.i./A) (including all application timings) to cotton per year under this application scenario.
- If environmental conditions prevent timely applications resulting in large weeds or heavy infestations, a single application of **Opportunity** at up to 43 fl. oz./A (0.80 lb. a.i./A) may be made to cotton. Do not make application of more than 43 fl. oz. of **Opportunity** in a single application under this use scenario. If a single application greater than 29 fl. oz. is made, a subsequent application not to exceed 29 fl. oz. may be made to cotton. The yearly total use rate under this scenario may not exceed 72 fl. oz. of **Opportunity** (1.33 lbs. a.i./A). Sequential applications need to be made at least 10 days apart.
- Do not make application of **Opportunity** through any type of irrigation system.

Refer to the **ROTATIONAL CROP RESTRICTIONS** section for the appropriate rotational crop information.

LibertyLink Cotton

Refer to the **WEED CONTROL FOR ROW CROPS** tables to select the proper application rate based upon the weeds present and their size. Uniform and thorough spray coverage is required to achieve consistent weed control. For ground application, make application of **Opportunity** to LibertyLink cotton as an over-the-top foliar spray directed to the lower one-third of the cotton stand.

Non-Glufosinate Tolerant Cotton

Application of **Opportunity** to cotton varieties not labeled as LibertyLink requires the use of hooded spray equipment designed to minimize exposure of the spray to the cotton stand. A hooded sprayer directs the spray onto weeds, while shielding the cotton stand from contact. Use nozzles that provide uniform coverage within the treated area. Keep hoods on these sprayers adjusted to protect desirable vegetation. Extreme care must be exercised to avoid exposure of the desirable vegetation to the spray.

With a hooded sprayer, the spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground. If the hoods are raised, spray particles may escape and come into contact with the cotton, causing damage or destruction of the crop.

Herbicide rates and spray volume instructions are presented as broadcast equivalents and must be reduced in proportion to the area actually treated. Use the following formulas to calculate the correct rate and volume per planted (field) acre:

<u>Band Width in Inches</u> Row Width in Inches	X	Broadcast Rate per Acre	=	Amount of Banded Product needed per Acre
<u>Band Width in Inches</u> Row Width in Inches	X	Broadcast Spray Volume per Acre	=	Banded Spray Volume needed per Acre

Post-Harvest

Opportunity may be applied as a post-harvest burndown treatment to fields (after cotton harvest). Up to 43 fl. oz./A of **Opportunity** may be applied in a single application to control larger weeds growing in the crop at the time of harvest. If more than 29 fl. oz./A is used in a single application, the yearly total may not exceed 72 fl. oz./A, including all application timings. Refer to the **ROTATIONAL CROP RESTRICTIONS** section for the appropriate rotational crop information.

Tank Mixtures - Cotton

Certain tank mixes may aid in the performance of **Opportunity**. No additional surfactant is needed with any tank mix partner. **Opportunity** may be applied in tank mix combination with labeled rates of other products provided these other products are labeled for the timing and method of application for the cotton to be treated. When tank mixing, do not exceed specified application rates. **Opportunity** cannot be mixed with any product containing a label prohibition against such mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

- **LibertyLink Cotton** - For cotton tolerant to **Opportunity**, Dual Magnum® or Staple® Herbicide may be tank mixed with **Opportunity** and applied over the top post-emergence to enhance weed control and/or provide residual control.
- **All Cotton Types** - The following herbicides may be tank mixed with **Opportunity** for hooded spray application to enhance weed control and/or provide residual weed control.

Post-Emergence Over-The-Top Tank Mix Partners for Opportunity on LibertyLink Cotton

Assure II	metolachlor	clethodim
Poast Plus	Fusilade DX	Select Max
Fusion	Staple	

DIRECTIONS FOR USE - SOYBEANS

Make application of **Opportunity** only to soybean designated as LibertyLink. Uniform, thorough spray coverage is necessary to achieve consistent weed control. For best results make application to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of **Opportunity**. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Adding ammonium sulfate with **Opportunity** may improve weed control if weeds are under stress. For optimal yield, early season weed removal is important.

Applications of **Opportunity** on soybeans may be made from emergence up to but not including the bloom growth stage. Make application of **Opportunity** to LibertyLink soybeans from emergence up to but not including the bloom growth stage at 29 to 43 fl. oz./A. See weed chart to determine rate. Should environmental conditions prevent a timely herbicide application, a single application of up to 43 fl. oz./A of **Opportunity** may be made to soybeans followed by one additional application at maximum of 43 fl. oz./A with a yearly maximum of 87 fl. oz./A. **Opportunity** may be applied alone or in a tank mix application with a residual herbicide to control weeds that have not yet emerged at the time of application.

Although timely post-applications of **Opportunity** can provide complete weed control, residual herbicides at burndown planting, or tank mixed with **Opportunity** help ensure optimal weed management, particularly if environmental conditions delay timely post-applications. Residual herbicides can also reduce early season weed competition and are a key element of good weed resistance management practices.

Use Pattern Rate Ranges		
1 st Application	2 nd Application	Yearly Maximum
29 - 43 fl. oz./A	29 - 43 fl. oz./A	87 fl. oz./A

Restrictions - Soybeans

- Do not make application of **Opportunity** within 70 days of harvesting soybean seed.
- Do not make application of more than 87 fl. oz./A (1.61 lbs. a.i./A) of **Opportunity** on soybeans per year.
- Do not make application of more than 43 fl. oz./A (0.80 lb. a.i./A) of **Opportunity** in a single application.
- Do not apply more than 2 applications per year.
- Do not graze the treated crop or cut for hay.
- Do not use nitrogen solutions as spray carriers. A silicone-based antifoam agent may be added if needed.
- Do not make application of **Opportunity** if soybeans show injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
- Sequential applications need to be at least 5 days apart.
- Do not make application of **Opportunity** through any type of irrigation system.

Refer to the **ROTATIONAL CROP RESTRICTIONS** section for the appropriate rotational crop information.

Tank Mixtures - Soybeans

Certain herbicide tank mixes may complement **Opportunity**. No additional surfactant is needed with any tank mix partner. **Opportunity** may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the soybean to be treated. When tank mixing, do not exceed specified application rates. **Opportunity** cannot be mixed with any product containing a label prohibition against such mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Tank Mix Partners for Opportunity in LibertyLink Soybeans

Assure® II	Fusion®	Raptor™
Classic®	Harmony® GT	Reflex®
clethodim	Optill	Resource®
Cobra®	metolachlor	Select Max®
Fierce	Phoenix™	Sharpen
FirstRate®	Poast Plus®	Synchrony® XP
Flexstar®	Prefix	Ultra Blazer®
Fusilade® DX	Pursuit®	

DIRECTIONS FOR USE ON CANOLA, CORN, COTTON, AND SOYBEAN SEED PROPAGATION

Opportunity may be applied to select out susceptible "segregates" (i.e., canola, corn, cotton, and soybean plants that are not tolerant to glufosinate-ammonium during seed propagation).

Canola: Opportunity may also be used in canola seed propagation as a foliar spray to selectively eliminate canola plants that do not carry a gene that imparts tolerance to glufosinate-ammonium and as such, can be applied to remove susceptible segregates during canola seed propagation. See **DIRECTIONS FOR USE - CANOLA** for use rates and application timing.

Corn: A hooded sprayer may be used to protect plants from coming into contact with the herbicide application. For the selection of tolerant corn segregates, **Opportunity** may be applied at 29 fl. oz./A plus AMS at 3 lbs./A (17 lbs./100 gals.) when corn is in the V-3 to V-4 stage of growth (i.e., 3 to 4 developed collars). A second treatment of 29 fl. oz./A plus AMS at 3 lbs./A may be applied when the corn is in the V-6 to V-7 stage of growth or up to 24" tall. Sequential applications need to be at least 7 days apart. When temperatures exceed 85°F, the rate of AMS can be reduced to 1.5 lbs./A (8.5 lbs./100 gals.) to reduce potential leaf burn.

Cotton: Opportunity may also be used in cotton seed propagation as a foliar spray to selectively eliminate cotton plants that do not carry a gene that imparts tolerance to glufosinate-ammonium and as such, can be applied to remove susceptible segregates during cotton seed propagation. See **DIRECTIONS FOR USE - COTTON** for use rates and application timing.

Soybean: For the selection of tolerant soybean (segregates), **Opportunity** may be applied at up to 29 to 43 fl. oz./A when soybean is in the third trifoliolate stage. A second treatment of 29 to 43 fl. oz./A may be applied up to but not including the bloom growth stage of soybean. Sequential applications need to be at least 5 days apart.

Restrictions

- **Canola & Cotton:** Breeding material not possessing the glufosinate-ammonium tolerance gene will be severely injured or killed if treated with this herbicide.
- **Corn:** Inbred lines (plants not possessing glufosinate-ammonium tolerance) will be severely injured or killed if treated with this herbicide.

DIRECTIONS FOR USE ON LISTED TREE, VINE, AND BERRY CROPS

Bushberries (blueberry, currant, elderberry, gooseberry, and huckleberry); Other Berries (lingonberry, juneberry, and salal); Citrus (lemon, orange, grapefruit, lime, mandarin, tangerine, tangelo, calamondin, kumquat, pummelo, citron, citrus hybrids, tangor, and cultivars, varieties and/or hybrids of these); Olives; Pome Fruit (apple, pear, crabapple, loquat, mayhaw, quince, azarole, medlar, tejocote, cultivars, varieties and/or hybrids of these); Stone Fruit (apricot, cherry, peach, nectarine, plum, capulin, jujube, sloe, and cultivars, varieties and/or hybrids of these); Tree Nuts (almonds, filberts, hickory nuts, macadamia nuts (bush nuts), pecans, pistachios, and walnuts); Vineyards (all grape varieties (table, wine, and raisins))

Make application of **Opportunity** to the tree, vine, and berry crops listed below. Uniform, thorough spray coverage is necessary to achieve consistent weed control. For best results, make application to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of **Opportunity**. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Weeds under stress or in dense populations will require application at the highest specified label use rate. Stressed conditions also include prior treatments of other contact or systemic herbicides. Do not retreat these weeds with **Opportunity** until sufficient regrowth has occurred.

Make application of **Opportunity** as a directed spray to control undesirable vegetation in tree, vine, and berries listed on this label. Make application as a broadcast, banded, or spot treatment application depending on the situation to control weeds listed under the **Weeds Controlled in Tree, Vine, and Berry Crops** table. Avoid direct spray or drift to desirable vegetation. Regrowth may occur due to the weed stage of growth at application, low use rate, or environmental conditions. Repeat applications of **Opportunity** may be necessary to control plants generating from underground parts or seed.

Avoid contact of **Opportunity** solution, spray, drift or mist with green bark, stems, or foliage, as injury may occur to trees, vines, and berries. **Only trunks with callused mature brown bark should be sprayed unless protected from spray contact by nonporous wraps, grow tubes, or waxed containers. Contact of Opportunity with parts of trees, vines, or berries other than mature brown bark can result in serious damage.**

Broadcast Applications

Make application of **Opportunity** at the rates listed below for broadcast applications based on weed size and stage of growth.

Weed Size and Stage	Opportunity Rate
Weeds < 3" in height	48 fl. oz./A (0.89 lb. a.i./A)
Weeds < 6" in height pre-tiller grasses	56 fl. oz./A (1.04 lbs. a.i./A)
Weeds > 6" in height and/or grasses that have tilled	56 - 82 fl. oz./A (1.04 - 1.52 lbs. a.i./A)

Banded Spray Applications

Banded applications may be used using the following formula to calculate the amount of herbicide needed for orchard or vineyard strip sprays:

$$\frac{\text{Band Width in Inches}}{\text{Row Width in Inches}} \times \text{Rate per Acre Broadcast} = \text{Amount of Herbicide needed for Treatment}$$

Spot or Directed Spray Applications

For spot or directed spray applications by backpack sprayers only (no mechanically pressured handgun applications allowed), mix **Opportunity** at 1.7 fl. oz. of product per gallon of water. Make application to undesirable vegetation foliage until wet but before runoff. Ensure uniform and complete coverage. Thoroughly clean the sprayer following use. **DO NOT** make spot or directed spray applications to tree or vine trunk as injury may occur.

Restrictions - Tree, Vine, and Berry Crops

- Do not make application of more than 164 fl. oz. of **Opportunity** per acre (3 lbs. a.i./A) to berry bushes and stone fruit in a 12 month period. Do not make more than 2 applications at a maximum rate of 82 fl. oz. per acre (1.5 lbs. a.i./A) per application.
- Do not make application of more than 246 fl. oz. (4.5 lbs. a.i./A) of **Opportunity** per acre to tree nuts, vines, pome fruit, citrus, and olives in any calendar year. Do not make more than 3 applications at a maximum rate of 82 fl. oz. per acre (1.5 lbs. a.i./A) per application.
- Do not graze, harvest, and/or feed treated orchard cover crops to livestock.
- Do not make application of this product aerially to tree, berry, or vine crops.
- Do not make application of this product within 14 days of nut, fruit, berry, or grape harvest.
- Applications to citrus fruits, pome fruits, and olives must be a minimum of 14 days apart.
- Applications to stone fruit must be a minimum of 28 days apart.
- Do not make spot spray applications to suckers, as tree injury may occur.
- Do not make application of **Opportunity** through any type of irrigation system.

Sucker Control with Opportunity

Opportunity will reduce or eliminate sucker growth when applied to suckers that are young, green, and uncalloused. For sucker control, make application of a split application approximately 4 weeks apart at 56 fl. oz. of **Opportunity** per acre. Coverage of all sucker foliage is necessary for optimum control. Suckers should not exceed 12" in length.

Tank Mixtures - Tree, Vine, and Berry Crops

Opportunity does not provide residual weed control or control of unexposed plant parts. Certain herbicide tank mixes may aid in the performance of **Opportunity** or be added to provide residual herbicide activity. No additional surfactant is needed with any tank mix partner. **Opportunity** may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the crop to be treated. When tank mixing, do not exceed specified application rates. **Opportunity** cannot be mixed with any product containing a label prohibition against such mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Chateau	Princep® 4L	Sinbar® 80W
Devnnot® 50WP	Simazine 4L	Solicam® DF
Goal® 1.6E	Simazine 80W	Surflan® A/S
Karmex® DF	Simazine 90	

Weeds Controlled in Tree, Vine, and Berry Crops

Broadleaf Weeds			
Alkali Sida	Fleabane, Annual	Morningglory, Ivyleaf	Smartweed, Pennsylvania
Ammannia, Purple	Goosefoot	Morningglory, Pitted	Sowthistle, Annual
Arrowhead, California	Gromwell, Field	Mullein, Turkey	Spurge, Prostrate
Buckwheat, Wild	Groundcherry, Cutleaf	Mustard, Wild	Starthistle, Yellow
Buffalobur	Groundsel, Common	Nettle	Sunflower, Common
Burclover, California	Henbit	Nightshade, Black	Sunflower, Prairie
Carpetweed	Jimsonweed	Nightshade, Eastern Black	Sunflower, Volunteer
Chickweed, Common	Knotweed	Nightshade, Hairy	Swinecress
Chinese, Thornapple	Kochia	Pennycress	Thistle, Russian
Cocklebur, Common	Lambsquarters, Common	Pigweed, Redroot	Turnip, Wild
Copperleaf, Virginia	Lettuce, Miner's	Pineapple Weed	Velvetleaf
Cudweed	Lettuce, Prickly	Puncturevine	Vervain
Cutleaf Evening Primrose	London Rocket	Purslane, Common	Vetch
Dodder	Mallow, Common	Radish, Wild	Willowherb, Panicle
Eclipta	Malva (Little Mallow)	Ragweed, Common	
Fiddleneck	Marestail	Ragweed, Giant	
Filaree	Mayweed	Redmaids	
Filaree, Redstem	Morningglory, Entireleaf	Shepherd's Purse	
Grass Weeds			
Barnyardgrass	Crabgrass, Smooth	Junglerice	Shattercane
Bluegrass, Annual	Cupgrass, Woolly	Oat, Wild	Sprangletop
Brome, Rippgut	Foxtail, Giant	Panicum, Fall	Stinkgrass
Bromegrass, Downy	Foxtail, Green	Panicum, Texas	Wheat, Volunteer
Canarygrass	Foxtail, Yellow	Rush, Toad**	Windgrass
Chess, Soft	Goosegrass	Ryegrass, Annual*	Witchgrass
Crabgrass, Large	Johnsongrass, Seedling	Sandbur, Field	
Biennial and Perennial Weeds			
Aster, White Heath	Dallisgrass	Mustard, Tansy	<i>Rubus</i> spp.
Bindweed, Field	Dandelion	Nutsedge, Purple	Spurge, Leafy
Bindweed, Hedge	Dock, Curly	Nutsedge, Yellow	Thistle, Bull
Bluegrass, Kentucky	Dogbank, Hemp	Onion, Wild	Thistle, Musk
Bromegrass, Smooth	Fescue	Orchardgrass	Torpedograss
Bulrush*	Goldenrod, Gray	Paragrass	Vaseygrass
Burdock	Guineagrass	Plantain	Woodsorrel
Canada Thistle	Horsetail	Poison Ivy/Oak	Yarrow, Common
Clover, Alsike	Love Grass	Quackgrass	
Clover, Red	Mugwort	Rocket, Yellow	
Clover, White	Mullein, Common	Rose, Wild	
*Make application to annual ryegrass before 3" in height.			
**Indicates suppression.			

DIRECTIONS FOR POTATO VINE DESICCATION

Make application of **Opportunity** at the beginning of natural senescence of potato vines. Make application of 21 fl. oz./A. Do not split this application or apply more than one application per harvest. Potato varieties with heavy or dense vines may require an application of another desiccation product to complete vine desiccation.

Thorough coverage of the potato vines to be desiccated is essential. Use a sufficient volume of water (20 to 100 gals./acre) to obtain a thorough coverage of the potato vines. Vary the gallons of water per acre and the spray pressure as indicated by the density of the potato vines to assure thorough spray coverage. Increase the spray volume to at least 30 gals. of water per acre when the potato vine canopy is dense or under cool and dry conditions. Make application of **Opportunity** with the spray boom as low as possible to achieve thorough coverage of the potato vines for best control and to minimize drift potential.

Restrictions - Potato Vine Desiccation

- Do not make application of more than 21 fl. oz./A (0.39 lb. a.i./A) of **Opportunity** to potato vines per year.
- Do not apply more than one application per year.
- Do not harvest potatoes until 9 days or more after application of **Opportunity**.
- Do not make application to potatoes grown for seed.
- Canola, corn, cotton, potatoes, soybean, and sugar beets may be planted at any time after the application of **Opportunity** as a potato vine desiccant.
- Do not plant treated areas to wheat, barley, buckwheat, millet, oats, rye, sorghum, and triticale until 30 or more days after an application of **Opportunity** as a potato vine desiccant.
- Do not plant treated areas to root and tuber vegetables, leafy vegetable, and brassica vegetable until 70 days after an application of **Opportunity** as a potato vine desiccant.
- Do not plant treated areas to crops other than those listed in this use precautions section until 120 or more days after an application of **Opportunity** as a potato vine desiccant.

FALLOW FIELDS OR POST-HARVEST

Opportunity may be used as a substitute for tillage in fallow fields to control or suppress weeds listed in the **WEED CONTROL FOR ROW CROPS** section. Applications may be made in fallow fields, post-harvest, before planting or emergence of any crop listed on this label.

Make application of **Opportunity** at 22 or 29 fl. oz./A to fallow fields to control specific weeds. **Opportunity** must be applied with ammonium sulfate. Tank mixes with 2,4-D, glyphosate or atrazine may be used with **Opportunity** to enhance total weed control. When tank mixing, do not exceed specified application rates. **Opportunity** cannot be mixed with any product containing a label prohibition against such mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. See the **APPLICATION AND MIXING PROCEDURES** section for additional information on how to apply this product. Refer to the **ROTATIONAL CROP RESTRICTIONS** section for the appropriate rotational crop information.

FARMSTEADS, RECREATIONAL, AND PUBLIC AREAS

When applied as listed, **Opportunity** controls undesirable plant vegetation in non-crop areas around farmstead, building foundations, shelter belts, along fences, airports, commercial plants, storage and lumber yards, educational facilities, fence lines, ditch banks, dry ditches, schools, parking lots, tank farms, pumping stations, parks, other public areas, and non-selective farmstead weed control. Refer to the **DIRECTIONS FOR USE - LISTED TREE, VINE, AND BERRY CROPS** section for appropriate application broadcast and spot spray application rates and lists of weeds controlled.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Do not use or store near heat or open flame. Keep the container tightly closed and dry in a cool, well ventilated place. Storage temperature should not exceed 125°F. Protect against direct sunlight.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING [Less Than 5 Gallons]: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONTAINER HANDLING [Greater Than 5 Gallons]: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

CONTAINER HANDLING [For Bulk and Mini-Bulk Containers]: Refillable container. Refill this container with pesticide only. Do not use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by State and local authorities.

SEED DISPOSAL: To dispose of out of date or otherwise unmarketable seed from plants which have been treated with this product, broadcast and lightly incorporate seed into field soils using disc or other suitable implement. Any resulting crop may be destroyed by chemical or mechanical means. Alternatively, seed may be destroyed by deep burial, incineration or landfill disposal.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, Sharda USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

All trademarks are the property of their respective owners.

Opportunity

GROUP 10 HERBICIDE

OPEN HERE

A non-selective herbicide for post-emergence broadcast use on canola, sweet corn*, corn, cotton, and soybean designated as LibertyLink®. Opportunity may be used for weed control in non-glufosinate tolerant cotton when applied with a hooded sprayer in-crop. Opportunity may also be applied as a broadcast burndown application before planting or before emergence of canola, sweet corn*, corn, soybean, and sugar beet* designated as LibertyLink® and any conventional canola, sweet corn*, corn, cotton, soybean, or sugar beet. Opportunity may be used for post-emergence weed control on olives, listed tree, vine, and berry crops. Opportunity may also be applied for potato vine desiccation.

*Not for use in California.

ACTIVE INGREDIENT:	WT. BY %
Glufosinate ammonium*	24.5%**
OTHER INGREDIENTS:	75.5%
TOTAL:	100.0%

*CAS Number 77182-82-2.

**Equivalent to 2.37 lbs. of active ingredient per U.S. gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

See label booklet for complete Precautionary Statements and Directions For Use.

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">Take off contaminated clothing.Rinse skin immediately with plenty of water for 15-20 minutes.Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none">Hold eye open and rinse slowly and gently with water for 15-20 minutes.Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none">Call a poison control center or doctor immediately for treatment advice.Have person sip a glass of water if able to swallow.Do not induce vomiting unless told to by a poison control center or doctor.Do not give anything to an unconscious person.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222.	
Note to Physician: If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible followed by charcoal and sodium sulfate administration.	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Do not use or store near heat or open flame. Keep the container tightly closed and dry in a cool, well ventilated place. Storage temperature should not exceed 125°F. Protect against direct sunlight.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING [Less Than 5 Gallons]: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinse into application equipment or mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONTAINER HANDLING [Greater Than 5 Gallons]: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

CONTAINER HANDLING [For Bulk and Mini-Bulk Containers]: Refillable container. Refill this container with pesticide only. Do not use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by State and local authorities.

SEED DISPOSAL: To dispose of out of date or otherwise unmarketable seed from plants which have been treated with this product, broadcast and lightly incorporate seed into field soils using disc or other suitable implement. Any resulting crop may be destroyed by chemical or mechanical means. Alternatively, seed may be destroyed by deep burial, incineration or landfill disposal.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

Manufactured For: Sharda USA, LLC, 7217 Lancaster Pike, Suite A, Hockessin, Delaware 19707
EPA Reg. No.: 83529-82 EPA Est. No.: 70815-GA-002, 39578-TX-001, 05905-IA-001, 228-IL-001 Net Contents: 2.5 Gallons